



2022

ABSTRACT BOOK

INTERNATIONAL SOIL SCIENCE SYMPOSIUM on SOIL SCIENCE & PLANT NUTRITION

(7th International Scientific Meeting)

2 – 3 December 2022

Samsun, Türkiye

Editors

Dr.Rıdvan KIZILKAYA

Dr.Coşkun GÜLSER

Dr.Orhan DENGİZ

Organized by

Federation of Eurasian Soil Science Societies

Erasmus Mundus Joint Master Degree in Soil Science (emiSS) Programme



With the support of the
Erasmus+ Programme
of the European Union

Cover design by FESSS

Editors:

Dr.Rıdvan Kızılkaya

Ondokuz Mayıs University, Faculty of Agriculture
Department of Soil Science and Plant Nutrition
55139 Samsun, Türkiye

Dr.Coşkun Gülser

Ondokuz Mayıs University, Faculty of Agriculture
Department of Soil Science and Plant Nutrition
55139 Samsun, Türkiye

Dr.Orhan Dengiz

Ondokuz Mayıs University, Faculty of Agriculture
Department of Soil Science and Plant Nutrition
55139 Samsun, Türkiye

Copyright © 2022 by Federation of Eurasian Soil Science Societies.

All rights reserved

ISBN 978-605-63090-7-6

This Abstract book has been prepared from different abstracts sent to the symposium secretary only by making some changes in the format. Scientific committee regret for any language and/or aim-scope.

All rights reserved. No parts of this publication may be reproduced, copied, transmitted, transcribed or stored in any form or by any means such as mechanical, electronic, magnetic, optical, chemical, manual or otherwise, without prior written permission from copyright owner.

Publication date : 2 December 2022



Visit the Symposium web site at
<http://www.fesss.org/>

E-mail: symposium@fesss.org



With the support of the
Erasmus + Programme
of the European Union

Sensitivity of soil organic matter indicators to long-term (<50 yrs.) application of mineral nitrogen in Cambisol in Serbia

Elmira SALJNIKOV ^{1,2,*}, Nikola KOKOVIĆ ¹, Vladan UGRENOVIĆ ¹, Aneta BUNTIĆ ¹, Tara GRUJIĆ ¹, Ljubomir ŽIVOTIĆ ³, Goran JAČIMOVIĆ ⁴

¹ Institute of Soil Science, Department of Pedology, Teodora Drajzera 7, 11000, Belgrade, Serbia

² Mitscherlich Academy for Soil Fertility (MITAK) GmbH, Prof.-Mitscherlich-Allee 1, 14641 Paulinenaue, Germany

³ Department of Pedology and Geology, Faculty of Agriculture, Belgrade University, Serbia

⁴ Faculty of Agriculture, University of Novi Sad, Novi Sad, Sq. Dositeja Obradovića 8, 21000 Novi Sad, Serbia

ABSTRACT

The goal of production of higher grain yield is predominantly achieved by higher doses of synthesised and mineral fertilizers. Intensive and long-term addition of mineral nitrogen, often adversely affect soil properties and ultimately the adequate ecological functioning of the soil and its sustainability. Labile organic matter fractions are most sensitive soil parameter. Their functional balance is controlled by soil microorganisms. They influence the ecological stability and biological productivity of ecosystems by participating in the biochemical transformation of mineral fertilizers and the synthesis of biologically active substances and nitrogen fixation. On the experimental station set 60 years ago by the Institute of Soil Science in Belgrade, Serbia, the effect of increasing doses of nitrogen fertilizer on biological indices of Cambisol on a bipolar crop rotation (wheat/corn) was studied. Soil basal respiration (BR), potentially mineralizable C and N (PMN and PMC), microbial biomass C and N (MBC and MBN) and light-fraction OM (LFN and LFC) on Cambisol were determined in five treatments with increasing doses of N fertilizer. The parameters studied were significantly affected by the long-term application of mineral fertilizer compared with both the control and the adjacent native soil. The highest amounts of nitrogen (N150) did not significantly differ from N120 and N90 for most of the parameters studied. Potentially mineralizable C represented the largest labile carbon pool, while microbial biomass N was the largest labile nitrogen pool. The mineralization rates for C and N were oppositely distributed over the seasons. The sensitivity index correlated with the amount of light-fraction OM. The results give a deeper insight into the behaviour and distribution of different pools of labile SOM in the agro-landscapes and can serve as a reliable basis for further research focused on zero soil degradation.

Key words:

Corresponding author : Elmira SALJNIKOV

E-mail: soils.saljnikov@gmail.com