

International Symposium



**Water
Resources
Management:
New Perspectives
and
Innovative
Practices**

PROCEEDINGS BOOK

<http://www.swarm.ni.ac.rs/symposium>



**September 23rd – 24th, 2021
Novi Sad, Serbia**



Co-funded by the
Erasmus+ Programme
of the European Union



International Symposium

Water Resources Management: New Perspectives and Innovative Practices

Symposium Proceedings

23-24 September 2021, Novi Sad, Serbia

<http://www.swarm.ni.ac.rs/symposium>

Published by:

University of Novi Sad, Faculty of Technical Sciences

Editors:

Maja Petrović
Milan Gocić

Cover design:

Ljiljana Jevremović

Printed by:

Futura Novi Sad

Press:

150 copies

ISBN 978-86-6022-367-0

Scientific Committee

Michael Tritthart, University of Natural Resources and Life Sciences, Vienna
Zakhar Maletskyi, Norwegian University of Life Sciences
Panagiotis Prinos, Aristotle University of Thessaloniki
Maria Mavrova, University of Architecture, Civil Engineering and Geodesy
Barbara Karleuša, University of Rijeka, Faculty of Civil Engineering
Maria Manuela Portela, University of Lisboa
Slaviša Trajković, University of Niš
Srdan Kolaković, University of Novi Sad
Emina Hadžić, University of Sarajevo
Mili Selimotić, Dzemal Bijedic University of Mostar
Jelena Đokić, University of Pristina in Kosovska Mitrovica
Predrag Stanojević, Academy of Applied Science of Kosovo and Metohija
Goran Sekulić, University of Montenegro

Organizing Committee

Daniel Wildt, University of Natural Resources and Life Sciences, Vienna
Susann Andersen, Norwegian University of Life Sciences
Charalampos Skoulikaris, Aristotle University of Thessaloniki
Petar Filkov, University of Architecture, Civil Engineering and Geodesy
Bojana Horvat, University of Rijeka, Faculty of Civil Engineering
Rodrigo Proença de Oliveira, University of Lisboa
Milan Gocić, University of Niš
Maja Petrović, University of Novi Sad
Hata Milišić, University of Sarajevo
Marko Čeček, Dzemal Bijedic University of Mostar
Gordana Milentijević, University of Pristina in Kosovska Mitrovica
Jelena Rajović, Academy of Applied Science of Kosovo and Metohija
Ivana Čipranić, University of Montenegro
Olivera Gavrilović, Public Water Management Company “VodeVojvodine”

PREFACE

The University of Novi Sad, Faculty of Technical Sciences hosted the **International Symposium "Water Resources Management: New Perspectives and Innovative Practices"** from 23rd-24th September 2021. The Symposium was a part of the activities realized under the CBHE KA2 project SWARM "Strengthening of master curricula in water resources management for the Western Balkans higher education institutions (HEIs) and stakeholders" co-funded by the Erasmus+ Programme of the European Union. SWARM project is implemented by a consortium made up of seven HEIs from the Western Balkan Region and six HEIs from Program Countries: University of Niš, University of Novi Sad, University of Priština in Kosovska Mitrovica, University of Montenegro, University of Sarajevo, Džemal Bijedić University of Mostar, Academy of Applied Sciences of Kosovo and Metohija, University of Natural Resources and Life Sciences from Vienna, Norwegian University of Life Sciences, Aristotle University of Thessaloniki, University of Architecture, Civil Engineering and Geodesy from Sofia, University of Rijeka, Faculty of Civil Engineering, Universidade de Lisboa, and Public Water Management Company "Vode Vojvodine".

The main objective of the SWARM project is the education of water management experts in the Western Balkan Region in accordance with the national and European Union policies. This objective is further broken down into the following specific objectives:

- To improve the level of competencies and skills in higher education institutions by developing new and innovative master programmes in the field of water resources management in line with the Bologna requirements and national accreditation standards.
- To design and implement new laboratories in Western Balkan HEIs, in cooperation with project partners from Program Countries.
- to develop and implement LLL courses for professionals in water sector in line with EU Water Framework Directive.

The International Symposium was an interdisciplinary forum where the research results and best practices in the field of Water Resources Management were shared with all SWARM fellows and stakeholders from the entire world. At the same time, the event was the place for the promotion of the SWARM results to a wider audience.

We would like to express our sincere thanks to all session chairs, keynote speakers, presenters, Organizing and Scientific committees, as well as to many others who contributed to the success of this Symposium.

We are confident that the solid foundation created by the SWARM project will continue to build up and strengthen our unique international network.

In Novi Sad, September 2021

Symposium Chairs

Maja Petrović

Milan Gocić

CONTENTS

Milan Gocić, Michael Tritthart, Panagiotis Prinos, Charalampos Skoulikaris, Jelena Đokić, Olivera Gavrilović, Jelena Vojvodić, Emina Hadžić, Mili Selimotić, Goran Sekulić, Jelena Rajović, Maja Petrović, Ljiljana Jevremović, Slaviša Trajković	
STRENGTHENING WATER RESOURCES MANAGEMENT IN THE WESTERN BALKANS	1
Daniel Wildt, Michael Tritthart	
IAHR YOUNG PROFESSIONALS NETWORK	11
Luis Angel Espinosa, Maria Manuela Portela	
PERSPECTIVES OF A CLIMATE CRISIS: HIGHER RISKS FROM GLOBAL TO A SMALL ISLAND ENVIRONMENT	15
Emina Hadžić, Hata Milišić, Suvada Šuvalija	
WATER RESOURCES MANAGEMENT IN URBAN AREAS	19
Milica Marković, Mladen Milanović, Slaviša Trajković	
WATER QUALITY EVALUATION IN BOVAN RESERVOIR FOR IRRIGATION PURPOSE	25
Charalampos Skoulikaris, Panagiotis Kotsalis	
SIMULATION OF UNGAUGED BASINS IN CLIMATE CHANGE CONDITIONS	31
Slavka Bogdanova, Petar Filkov	
MULTI CRITERIA ANALYSIS FOR PRIORITIZATION OF INVESTMENTS FOR RECONSTRUCTION AND MODERNIZATION OF IRRIGATION INFRASTRUCTURE	37
Ivana Sušanj Čule, Barbara Karleuša, Bojana Horvat, Nevenka Ožanić	
MASTER'S THESIS IN THE FIELD OF HYDROTECHNICAL ENGINEERING AT THE FACULTY OF CIVIL ENGINEERING (UNIVERSITY OF RIJEKA) – GOOD PRACTICE EXAMPLES	45
Elvis Žic, Nevenka Ožanić	
THE FUNDAMENTALS OF RISK ASSESSMENTS ON THE GEOHAZARD CONSEQUENCES	51
Zorica Filipović, Ivana Lukić, Mladen Milanović	
HORIZONTAL LEGISLATION IN ENVIRONMENTAL PROTECTION, CASE STUDY: STUDENT PARTICIPATION IN MAKING A DECISION FOR WATER RESOURCE MANAGEMENT	57

Irma Dervišević, Almin Dervišević, Milica Tomović, Jovana Galjak	
WATER QUALITY ASSESSMENT OF RURAL WATER SUPPLIES OTHERWISE AND AFTER THE FLOOD ON THE TERRITORY OF THE CITY OF KRALJEVO AND THE MUNICIPALITY OF VRANJACKA BANJA	63
Davut Lacin, Vesna Teofilović, Sezen Kucuk, Jelena Pavličević, Ayse Z. Aroguz	
WASTEWATER TREATMENT USING HALLOYSITE/BIOPOLYMER NANOCOMPOSITE	69
Elvis Žic, Nevenka Ožanić	
OVERVIEW OF INPUT DATA FOR QUANTITATIVE RISK ANALYSIS FROM THE CONSEQUENCES OF GEOHAZARD	75
Miljan Jeremić, Milan Gocić	
VISUALIZATION OF AVERAGE ANNUAL PRECIPITATION IN SERBIA FOR THE PERIOD FROM 1946 TO 2019	81
Mladenka Novaković, Ivana Mihajlović, Goran Štrbac, Dragana Štrbac, Maja Petrović	
SUITABILITY ASSESSMENT OF PHOTOCATALYTIC TREATMENT FOR PHARMACEUTICAL REMOVAL – STRENGTH, WEAKNESS, OPPORTUNITIES AND THREATS (SWOT) ANALYSIS	89
Milena Ostojić, Ivana Čipranić, Goran Sekulić	
POSSIBILITIES OF APPLICATION OF HEC RAS TWO-DIMENSIONAL MODELS FOR PREDICTION OF BRIDGE PIER SCOUR	95
Danijela Milanović, Borislava Blagojević, Ljiljana Vasilevska	
NATURE BASED STORMWATER MANAGEMENT SOLUTIONS FOR HOUSING AREA – CASE STUDY OF ROOF GARDEN IMPLEMENTATION	101
Marija Perović, Vesna Obradović, Branislava Matić, Dragica Vulić	
TRACING THE NITROGEN SOURCE IN GROUNDWATER	107
Branislava Matić, Marija Perović, Dragica Vulić	
NATURAL WATER RETENTION MEASURES CONTRIBUTION TO INTEGRATED TRANSBOUNDARY TISZA RIVER BASIN MANAGEMENT-ENVIRONMENTAL AND FLOOD RISK MANAGEMENT OBJECTIVES SYNERGY	113

TRACING THE NITROGEN SOURCE IN GROUNDWATER

Marija Perović, Vesna Obradović*, Branislava Matić*, Dragica Vulić**

* Jaroslav Černi, Water Institute Jaroslava Černog 80, 11226 Belgrade

Abstract: The knowledge of physico-chemical and microbiological composition of groundwater indicates its usability and required conditioning treatments, affecting, and determining the value and price of water. The importance of maintaining the quality and quantity of groundwater is obvious from the fact that more than 97% of the total quantities of fresh water on Earth is groundwater. Due to the slow groundwater flow, pollutants reaching groundwater persist for a long time, requiring considerable efforts and financial investments to restore original water quality. In the case of nitrogen inflow in groundwater, the prevailing conditions will determine its fate. The aim of the research emphasizes the necessity of comprehensive research including geology, groundwater recharge regime, physico-chemical composition, to decrease uncertainties of the nitrogen origin determination by a single point of view.

Keywords: groundwater, isotopes, nitrogen

INTRODUCTION

Typical groundwater flow rates vary from 1 m/day to 1 m/year. Caused by the relatively slow groundwater velocity, comprehensive, complex, and specific processes of sorption, dissolution, convection, dispersion, and oxidation-reduction within the self-purifying potential of an aquifer are present (Dimkic et al., 2008). Unfolding of these processes usually leads to a significant improvement in water quality. Groundwater composition is conditioned by many factors: autochthonous (geological - aquifer type and genesis, recharge water quantity and quality) and allochthonous (sewage, industrial and agricultural pollution). Beside the difference in groundwater quality of different regions, the groundwater composition in terms of quality and quantity, may vary between shallow and deeper layers of the same aquifer.

About 75% of the EU population uses groundwater for water supply. In Serbia the share of groundwater used for water supply is 75% and it indicates the importance of maintaining the groundwater quality. Over 50% of the groundwater used for water supply in Serbia comes from alluvial groundwater sources. The soil fertility and availability of irrigation water made alluvial zones suitable for agricultural production. For groundwater management, it is important to consider the value of the aquifer's self-purifying potential, which in the case of agrochemicals (fertilizers and pesticides) may be particularly significant (Živančev et al., 2020).

Crop yields and fertilizer application are in a strong positive correlation. Fertilizer application, along with the productivity factors such as selection of crop varieties, irrigation, mechanization, and pesticide application have an obvious reflection in crop yield trends across the world in the 20th century. The groundwater contamination by fertilizers may last for many decades. The importance of agricultural nitrogen emission to the environment is reflected through "Nitrate Directive "(ND). In the Annex II of ND, a Code of good agricultural practice with the objective of reducing water pollution by nitrates covers the following items: periods when the land application of fertilizer is inappropriate; the land application of fertilizer to steeply sloping ground; the land application of fertilizer to water-saturated, flooded, frozen or snow-covered ground; the conditions for land application of fertilizer near water courses; the capacity and construction of storage vessels for livestock manures, including measures to prevent water pollution by run-off and seepage into the groundwater and surface water of liquids containing livestock manures and effluents from stored plant materials such as silage; procedures for the land application, including rate and uniformity of spreading, of both chemical fertilizer and livestock manure, that will maintain nutrient losses to water at an acceptable level (Directive 91/676/EEC, 1991). ND adopted by all EU Member States, limits the concentration of nitrate to 50 mg/l in all freshwater bodies (Directive 91/676/EEC, 1991). The prescribed nitrate concentration in drinking water and groundwater in Serbia is also 50 mg NO₃/l ("Official Gazette SRJ", br. 42/98 i 44/99 i " Official Gazette RS", br. 28/2019, " Official Gazette RS", br. 50/2012).

Nitrogen transformations are mostly conditioned by oxygen content, concentration, and bioavailability of electron acceptors (and donors), the ratio of electron donor and acceptors, dominant microbial population, pH value etc. Geology determines initial groundwater oxygen content. Oxygen content dominantly determines whether the nitrogen will remain in the most stable form (N^{+5}) or it will be reduced and lost from groundwater (N_2 , N_2O). If the concentration of dissolved oxygen is above 0.5 mg/l (and Eh > 250 mV) the nitrogen will probably remain in it's the most stable form (NO_3^-) which is unable to sorb on sediment particles. If there is lack of oxygen, the ratio of total organic carbon (TOC) to nitrates (and sulphide presence) will determine whether the nitrate reduction will lead to N loss (N_2 , N_2O) or conservation (NH_4^+) (Dissimilatory nitrate reduction to ammonium - DNRA/ Assimilatory Nitrate Reduction to ammonium - ANRA). The processes and conditions for nitrogen transformation are explained in detail in Perović et al., 2017; Perović et al., 2020.

MATERIALS AND METHODS

The competitive and often overlapping oxido-reducing reactions in groundwater induce difficulties in determination of the N origin considering only one scientific point of view. Determination of N origin and discrimination of conditions suitable for N conservation or loss, are of great importance in groundwater management. Although many factors affect nitrogen behaviour and fate in groundwater, and there is a wide range of nitrogen sources, the oxygen content, the boron concentration, and isotopic signatures of NO_3^- and NH_4^+ are selected for further investigation.

Oxygen in groundwater is predominantly used for oxidation of organic substances and lower oxidation state minerals (mainly iron), which are sorbed to fine-grained materials. The deposition of fine-grained materials is less pronounced in upper part of river basins due to a higher water flow energy. Therefore, the oxygen content is generally significantly higher in aquifers in the upstream parts of a basin (coarse-grained material), whereas anoxic aquifers are found in lower river basins (Figure 1) (Dimkić, 2012; Perović, 2019). The coarse-grained aquifers, with significant oxygen diffusion, or aquifers with intensive water exchange, are considered as sensitive to nitrate contamination.

If the aquifer is formed in lower part of the stream where the river flow is decreased, the river deposited the fine-grained material which is rich in iron and organic carbon, inducing anoxic conditions groundwater (Perović 2019). The anoxic aquifers are rich in oxygen consuming substances, mostly reduced metal forms and organic carbon.




Figure 1 Schematic representation of an alluvial aquifer (Dimkić, 2012)

RESULTS AND DISCUSSION:

Mineralization of organic matter is accompanied by a decrease in pH and often consequent reductive dissolution of Fe^{3+} ions (iron oxide and/or iron oxyhydroxide). Increased acidity of the environment causes the appearance of Fe^{2+} cations and often the release of the mobile form of arsenic, the arsenite cation (As^{3+}), which is a common constituent of anoxic groundwater in layers rich in arsenopyrite or iron-arsenite ores (Berg et al., 2001; Wang et al., 2017). If the medium is rich in organic matter, mineralization can generate the release of mobile ions Fe^{2+} , NH_4^+ As^{3+} when the observed linear correlation mainly indicates the autochthonous origin of ammonium cations.

As indicators of groundwater origin and anthropogenic impacts on its quality (in terms of nitrogen pollution), the chloride (Cl), sodium (Na) and boron (B), along with the isotopic signatures of ammonia and nitrates are usually quantified.

Natural origin of B may indicate seawater intrusion, may originate from dissolving sandstones and igneous rocks, or may be found in certain evaporates such as borax ($(\text{Na}_2\text{B}_4\text{O}_5[\text{OH}]_4 \cdot 8\text{H}_2\text{O})$) (Clark, 2015). Simultaneously elevated concentrations of B and NH_4^+ may indicate their common origin from sewage (Lindebaum, 2012). One of the most common anthropogenic uses of B compounds is the production of sodium perborate ($\text{NaBO}_3 \cdot n\text{H}_2\text{O}$), which is the component of whitening agents, in detergents, toothpastes and soaps. In liquid manure and septic tanks, B concentrations are significantly elevated (Nikolenko et al., 2018). High concentration of B followed by low concentration of NO_3^- is probably an indicator of anthropogenic influence (detergents), while moderate concentration of B followed by moderate concentration of NO_3^- is likely to indicate the influence of wastewater excrements (Widory et al., 2004).

One of the widely used approaches over the past few decades in nitrogen origin determination has involved stable isotope analysis. The heterogeneity of geochemical conditions, groundwater flow, wide range of nitrogen sources and agricultural practices, causes a significant range of the isotopic signature $\delta^{15}\text{N}-\text{NO}_3^-$. In groundwater below agricultural areas the isotopic signature $\delta^{15}\text{N}-\text{NO}_3^-$ varies from -8.3 to + 65.5 ‰ (Nikolenko et al., 2018). N originating from different sources is characterized by a different range of $\delta^{15}\text{N}-\text{NO}_3^-$ as well as different enrichment factor, which can be used to determine the origin and relative contribution of NO_3^- source to its content in groundwater (Nikolenko et al., 2018). In groundwater, due to fractionation during physico-chemical or biological reactions (NH_3 volatilization, nitrification, denitrification), the isotopic composition changes. Thus, the isotopic signature cannot be unambiguous in nitrogen origin determination. The lowest values of $\delta^{15}\text{N}-\text{NO}_3^-$ characterize mineral fertilizers, oxidation of soil organic matter, rainwater, respectively, while the highest values characterize manure or household wastewater, whose ranges may overlap (Table 1).

The range of data on the $\delta^{15}\text{N}-\text{NH}_4^+$ distribution in groundwater below agricultural areas is significantly more modest comparing to nitrates data. Published research have shown that the range of $\delta^{15}\text{N}-\text{NH}_4^+$ in aquifers is - 8.5 to + 23.8 ‰ (Kendall 1998; Nikolenko et al., 2018). Organic fertilizers, manure, sewage, and wastewater from treatment plants are generally considered to be the main anthropogenic sources of NH_4^+ ions in groundwater below agricultural land (Nikolenko et al., 2018). The lowest values of $\delta^{15}\text{N}-\text{NH}_4^+$ characterize rainwater, while the highest values were observed for manure and sewage wastewater. Organic matter shows slightly higher values of $\delta^{15}\text{N}-\text{NH}_4^+$ isotopic composition compared to mineral fertilizers and rainwater (Nikolenko et al., 2018). Known to date research results suggest that isotopic signatures from different NH_4^+ sources may overlap due to the specificity of the environment in certain areas (Figure 2).




Figure 2 Characteristic ranges of isotopic signatures of stable N isotopes depending on source (Perović, 2019; Perović and Dimkić, 2021)

Table 1 Transformation processes, necessary conditions and expected fractionation range (Perović, 2019)

PROCESS	CHEMICAL REACTION	NECESSARY CONDITIONS	FRACTIONATION (APPROXIMATELY,)
Fixation	$N_{2(g)} + 3 H_2O_{(g)} \rightarrow 2 NH_3 + 3/2 O_{2(g)}$	Lightning, Plant enzymes	0‰
Mineralization	$R - CHNH_2 - COOH + O \rightarrow R - CO - COOH + NH_3$	Decreasing pH, anoxic conditions until nitrification	$0 \pm 1‰$
Nitrification	$NH_4^+ + 1,5 O_2 \rightarrow NO_2^- + H_2O + 2H^+$ $NO_2^- + 0,5 O_2 \rightarrow NO_3^-$	pH decreasing, $O_2 > 1$ mg/l	-18 to -29 ‰ to -35‰ produced NO_3^- will be depleted in $\delta^{15}N$
Denitrification	$5CH_2O + 4NO_3^- \rightarrow 2N_2 + 4HCO_3^- + CO_2 + 3H_2O$	$O_2 < 2,0$ mg/l and $Eh < 250$ mV; pH increase, high TOC, non-sulphidic, low C:N; low TOC, low Fe, high C:N	by 5-40‰ enrichment of remaining nitrate
	$10 Fe^{2+} + 2 NO_3^- + 14 H_2O \rightarrow 10 FeOOH + N_2 + 18H^+$	Low TOC, high Fe,	
	$5HS^- + 8NO_3^- + 3H^+ \rightarrow 5SO_4^{2-} + 4N_2 + 4H_2O$	Sulphidic, FeS , S^0 high TOC	
Dissimilatory nitrate reduction to ammonium	$2H^+ + NO_3^- + 2CH_2O \rightarrow NH_4^+ + 2CO_2 + H_2O$	High TOC, non-sulphidic, high C:N-fermentative DNRA;	Remaining nitrate approximately enriches by +10‰, lighter NH_4^+ is produced
	$4HS^- + 4NO_3^- + 4H_2O + 4H^+ \rightarrow 4SO_4^{2-} + 4NH_4^+$	sulphidic, H_2S , high TOC,	
Anaerobic ammonium oxidation	$NH_4^+ + NO_2^- \rightarrow N_2 + 2H_2O$	Low TOC, low Fe, low C:N ($N > C$), respiratory DN to NO_2^- then anammox	Like denitrification

*Eh – electrochemical potential

CONCLUSION

Nitrate pollution of shallow aquifers in Europe and in the world is a common, important, and well-studied problem. Beside the application of nitrogen-based mineral fertilizers, unregulated sewage discharges as well as sewage leaks, the existence of septic tanks and manure applications are the most significant sources of elevated concentrations of nitrogen compounds in groundwater. Considering the nitrogen loadings, to maintain the basic, autochthonous groundwater quality and to determine the vulnerability of groundwater to certain pollution the simultaneous complex analysis of agrotechnical, physico-chemical and hydrogeochemical data must be conducted. To trace the sources of contaminants, beside the groundwater recharge and groundwater flow, prevailing conditions of the environment and their impact on specific pollutant transformation must be considered.

In conducted research it is suggested that nitrogen sources in groundwater under agricultural areas should be studied by simultaneous analysis of concentrations levels of O_2 , NH_4^+ , NO_3^- , TOC; changes in concentration levels of anthropogenic tracers B, Cl, Na; by examination of state condition parameters (pH, redox potential) and isotopic signatures of stable isotopes of nitrogen and oxygen. Beside physico-chemical analysis groundwater flow and recharge should also be considered. Analysis of the ratio of stable isotopes of nitrogen (isotopic signatures) is a widely used method for identification of nitrate source in groundwater. As the isotopic signature of nitrogen can change under the influence of various complex biogeochemical mechanisms of transformation it cannot be exclusive in the source identification. Verification of the assumed transformation processes and nitrogen sources in groundwater is required by simultaneous, comprehensive, and complex analysis of physico-chemical, hydrogeological, and isotopic signatures.

ACKNOWLEDGMENT

The work was supported by Ministry of Education, Science and Technology Development of the Republic of Serbia under the Project “Methodology for the Assessment, Design and Maintenance of Groundwater Sources in Alluvial Environments Depending on the Aerobic State”, No. TR37014.

REFERENCES

- Berg M., Tran H.C., Nguyen T.C., Pham H.V., Schertenleib R., Giger W. 2001 Arsenic contamination of groundwater and drinking water in Vietnam: A human health threat. *Environmental Science & Technology*, 35(13), 2621-2626.
- Clark I. 2015 Contaminant Geochemistry and Isotopes. In: *Groundwater Geochemistry and Isotopes*. Taylor & Francis Group, LLC, Boca Raton, FL.
- Dimkić M. A., Brauch H.-J., Kavanaugh M. 2008 Groundwater management in large river basins, IWA Publishing, Alliance House, 12 Caxton Street, London SW1H 0QS, UK.
- Dimkić M. 2012 Importance of the Aerobic State of Alluvial Aquifers for Groundwater Use, 2012 Nakdong River International Water Week, 7th World Water Forum, 12-13 September 2012, Andong, Republic of Korea, pp 69-96.
- Kendall C. 1998 Tracing nitrogen sources and cycling in catchments, In: Kendall, C., McDonnell J.J. Eds., *Isotope Tracers in Catchment Hydrology*, Elsevier, Amsterdam, 519-576.
- Lindebaum J. 2012 Identification of sources of ammonium in groundwater using stable nitrogen and boron isotopes in Nam Du, Hanoi. PhD Thesis, Geology at Lund University, Sweden.
- Nikolenko O., Jurado A., Borges A. V., Knöller K., & Brouyre S. 2018 Isotopic composition of nitrogen species in groundwater under agricultural areas: a review. *Science of the Total Environment*, 621, 1415–1432.
- Perović M., Obradović V., Kovačević S., Mitrović D., Živančev N., & Nenin T. 2017 Indicators of groundwater potential for nitrate transformation in a reductive environment. *Water Environment Research*, 89(1), 4–16.
- Perović M. 2019. Procena uticaja i regionalne specifičnosti hidrogeohemijskih uslova na transformaciju azotnih jedinjenja u podzemnim vodama (Impact assessment and regional specificities of hydrogeochemical

conditions on the transformation of nitrogenous compounds in groundwater) PhD Thesis, University of Novi Sad Serbia.

Perović M., Obradović V., Zuber-Radenković V., Obrovski B., Miloradov-Vojinović M., Dimkić M. 2020 Comprehensive Biogeochemical Analysis of Nitrogen Transformation Parameters. *Water Resources*, 47(1), 156-170.

Perović M., and Dimkić M. 2021 Chapter 8 Transformation of nitrogen compounds in groundwater. In: *Alluvial Aquifer Processes*, IWA Publishing, Alliance House, 12 Caxton Street, London ISBN13:9781789060898.

Weng T. N., Liu C. W., Kao Y.H., Hsiao S.S.Y. 2017 Isotopic evidence of nitrogen sources and nitrogen transformation in arsenic-contaminated groundwater. *Science of Total Environment*, 578 167-185.

Widory D., Kloppmann W., Chery L., Bonnin J., Rochdi H., Guinamant J.L. 2004 Nitrate in groundwater: an isotopic multi-tracer approach, *J. Contam. Hydrol.* 72, 165–188.

Živančev N., Kovačević S., Perović M., Čalenić A., Dimkić M. 2020 Influence of oxic and anoxic groundwater conditions on occurrence of selected agrochemicals. *Water Supply*, 20(2) ISSN 1606-9749 EISSN 1607-0798

ORGANIZER & PUBLISHER

University of Novi Sad
Faculty of Technical Sciences



SUPPORTER



Co-funded by the
Erasmus+ Programme
of the European Union





Co-funded by the
Erasmus+ Programme
of the European Union



CERTIFICATE OF ATTENDANCE

This is to certify that

Marija Perović

ATTENDED THE

Symposium

Water Resources Management: New Perspectives and Innovative Practices

September 23rd - 24th, 2021 | Novi Sad | Serbia

A handwritten signature in black ink, appearing to read 'M. Petrović', written over a horizontal line.

Dr Maja Petrović
Symposium Chair
Faculty of Technical Science
University of Novi Sad

A handwritten signature in black ink, appearing to read 'M. Gocić', written over a horizontal line.

Dr Milan Gocić
Symposium Co-chair
Faculty of Civil Engineering and Architecture
University of Niš