



VII EUROSOIL 2025
& X Congreso Ibérico
de la Ciencia del Suelo

SEVILLE-SPAIN 8-12 SEP

**Advancing
soil knowledge for
a sustainable future**
Book of abstracts



VII EUROSOIL 2025 & X Congreso Ibérico de la Ciencia del Suelo

SEVILLE-SPAIN 8-12 SEP



ORGANIZED BY:



Advancing Soil Knowledge for a Sustainable Future

Book of Abstracts
of the Communications
presented to the
VII EUROSOIL Meeting
Seville – Spain
September 8 – 12, 2025

Title: Advancing Soil Knowledge for a Sustainable Future – EUROSOIL 2025 Book of Abstracts*

Editors: José A. González Pérez & José María de la Rosa Arranz

Published on-line in:

Digital.CSIC (<http://digital.csic.es/>), the Institutional Repository of "Consejo Superior de Investigaciones Científicas" (CSIC).

© VII EUROSOIL, Seville, Spain 2025

URI: <http://hdl.handle.net/10261/398890>

ISBN: 978-84-09-75471-7

Editors:

José A. González Pérez and José María de la Rosa Arranz



VII EUROSOIL 2025 & X Congreso Ibérico de la Ciencia del Suelo

SEVILLE-SPAIN 8-12 SEP



ORGANIZED BY:



CONTENT

PRESENTATION OF THE VII EUROSOIL CONGRESS	7
ORGANIZING COMMITTEE	8
SCIENTIFIC COMMITTEE	9
SEVILLA, VENUE & HOW TO GET THERE?	10
INVITED SPEAKERS	12
EUROSOIL 2025 GENERAL TOPICS	16
SPECIAL EVENTS WITHIN EUROSOIL 2025	23
EUROSOIL 2025 WORKSHOPS	24
EUROSOIL 2025 ASSOCIATIONS ASSEMBLIES	26
SOCIAL EVENTS WITHIN EUROSOIL 2025	26
EUROSOIL 2025 TECHNICAL SECRETARIAT	26
SPONSORS	27
SCIENTIFIC PROGRAMME	29
SUMMARY OF THE DAILY PROGRAM	91
PLENARY CONFERENCES	93
PLENARY CONFERENCES	94

ABSTRACTS OF THE COMMUNICATIONS

EU Soil Observatory (EUSO) 5 th Stakeholders Forum	101
GT 01. Soil Genesis, Evolution and Classification	133
GT 02. Soil Physical Chemistry	177
GT 03. Soil Biogeochemistry & Global Change	211
GT 04. Soil Biology 2	293
GT 05. Soil Hydrology & Erosion	377
GT 06. Soil Productive Sectors Economy and Sustainability	421
GT 07. Soil Amendments & Fertilizers	437
GT 08. Soil-Plant-Water Interactions	611
GT 09. Soil Health	671
GT 10. Soil Carbon Dynamics & Stabilization	793
GT 11. Soil Degradation, Desertification, Contamination - Conservation, Mitigation, Remediation	915
GT 12. Soil Natural Hazards Fires, Floodings, Landslides	1047
GT 13. Land Fragmentation, Urban and Anthropogenic Soils-POSTER-MARK	1067
GT 14. Soil Data Acquisition, Curation, Sharing & Modelling	1093
Gt 16. Soil Literacy Education & Art	1171
GT 17. Soils At The Crossroads Society, Policies, And Participatory Science	1197
AUTHORS INDEX	1239



VII EUROSOIL 2025 & X Congreso Ibérico de la Ciencia del Suelo

SEVILLE-SPAIN 8-12 SEP

GT 09 - 687 - P

IMPACT OF CEREAL/LEGUME INTERCROPPING SYSTEM ON MYCORRHIZAL ABUNDANCE AND ITS INTERACTION WITH C SEQUESTRATION IN THE ORGANICALLY MANAGED CROPPING SYSTEM

GT 09. SOIL HEALTH / GT 10. SOIL CARBON DYNAMICS AND STABILIZATION

AGNE VERŠULIENE*^{1,2}, ANDRIUS GARBARAS¹, MONIKA TOLEIKIENE²

¹ Center for Physical Sciences and Technology, Vilnius, Lithuania

² Lithuanian Research Centre for Agriculture and Forestry, Kedainiai, Lithuania

* agne.versulienė@ftmc.lt

Arbuscular mycorrhizal (AM) fungi form mutualistic relationships with the majority of land plants and are an important part of the soil microbial community in natural and agricultural ecosystems. These fungi promote water and nutrient acquisition by their host plant and regulate the allocation of photosynthetic carbon to soil. Both crop variety and environment affect naturally occurring mycorrhizal abundance in roots, but the relative importance of those factors for mycorrhization is largely unknown. Therefore, this study aims to investigate the impact of cereal/legume intercropping systems on mycorrhizal abundance and C sequestration in the organically managed cropping system.

The field study experiment was conducted in the Lithuanian Research Centre for Agriculture and Forestry, Akademija, Kedainiai district (55°23'50"N, 23°51'40"E) in the organically managed cropping system in 6 treatments: two monocultures (spring barley and oat) and two compositions with legume intercropping (field pea and red clover). Mycorrhizal colonisation intensity (M%) of the root sample segments was determined by light microscopy (Trinchera et al., 2019) in the LAMMC Microbiology Laboratory. The potential of the C sequestration in the system was estimated by incubating the intercropping systems using the ¹³C-CO₂ pulse labelling method (De Neergaard and Gorissen, 2004). The content of carbon isotope in plant roots and soil was assessed using a Thermo Flash EA 1112 elemental analyser interfaced to a Thermo Scientific Delta V Advantage isotope ratio mass spectrometer (IRMS) in the Isotopic Research Laboratory at the Centre for Physical Sciences and Technology.

Spring barley and oats M% ranged from 29 to 44% and 44 to 63%, respectively, across all samples. Averaged across all samples, the M% of oat was significantly higher than spring barley. The establishment of field pea intercrop has resulted in a significantly higher M% on spring barley and tended to decrease on oat. The red clover intercrop showed a positive effect on increasing M% on oat. The M% of spring barley and oat positively correlated with yield and showed significant differences in root biomass and C content in the soil.

ACKNOWLEDGEMENTS

This project has received funding from the Research Council of Lithuania (LMTLT), agreement No S-PD-24-39.

REFERENCES

de Neergaard A., Gorissen A. 2004. Carbon allocation to roots, rhizodeposits and soil after pulse labelling: a comparison of white clover (*Trifolium repens* L.) and perennial ryegrass (*Lolium perenne* L.). *Biology and Fertility of Soils*, 39: 228–234. doi:10.1016/j.scitotenv.2021.150857.

Trinchera A., Ciaccia C, Testani E., Baratella V., Campanelli G., Leteo F., Canali S. 2019. Mycorrhiza-mediated interference between cover crop and weed in organic winter cereal agroecosystems: The mycorrhizal colonization intensity indicator. *Ecology and Evolution*, 1-12. <https://doi.org/10.1002/ece3.5125>

Veršulienė A., Hirte J., Ciulla F., Camenzind M., Don A., Durand-Maniclas F., Heinemann H., Herrera J. M., Hund A, Seidel F., da Silva-Lopes M., Toleikienė M., Visse-Mansiaux M., Yu K., Bender S. F. 2024. Wheat varieties show consistent differences in root colonization by mycorrhiza across a European pedoclimatic gradient. *European Journal of Soil Science*, 75(4). <https://doi.org/10.1111/ejss.13543>



ORGANIZED BY:



INDEX

GT 09 - 696 - P

TRACE ELEMENTS IN THE SOIL-PLANT SYSTEM: ALFALFA AND NICKEL CASE STUDY

GT 09. SOIL HEALTH / GT 07. SOIL AMENDMENTS & FERTILIZERS

MIRA MILINKOVIC*¹, VESNA MRVIC¹, MILA PEŠIĆ¹, SONJA TOŠIĆ JOJEVIĆ¹, SNEŽANA ANDJELKOVIĆ², BILJANA SIKIRIĆ¹, OLIVERA STAJKOVIC-SRBINOVIC¹

¹ Institute of Soil Science, Belgrade, Serbia

² Institute for Forage Crops, Kruševac, Serbia

* miramilinkovic@yahoo.com

In Serbian soils, nickel (Ni) and chromium (Cr) are found in elevated concentrations over significant areas, often these elements origin is geochemical (from basic and ultrabasic rocks which account for about 4% of territory), but it can also be the result of anthropogenic pollution. Elevated Ni and Cr concentrations often occur in soils, which are generally suitable for agricultural production, including alfalfa cultivation. With the aim to select the nickel contaminated soil, agricultural soils from 15 locations in the central part of the country were examined for basic soil properties, and content and solubility of heavy metals. The soils with contrasting Ni concentrations (with slightly elevated and remediation Ni concentrations) were selected for further monitoring of alfalfa-soil interactions and elements translocation. The both soils were slightly acidic, moderately supplied with humus, nitrogen and available potassium, with clay mechanical composition. In addition, the inoculation with effective nitrogen-fixing bacteria rhizobia was performed at the beginning of trial, to ensure stable yields, and to examine the effect of strains on the accumulation of trace metals in the above-ground part of plants.

Lower solubility of metals in weakly acidic soil and adsorption on clay minerals possibly affected the lower mobility and accessibility of elements in the soil.

There is a pronounced accumulation of elements in the roots of the plant, which is confirmed by the translocation factor, which is generally less than 1. The Ni concentrations in alfalfa shoots varied depending on the strain, soil Ni and growing conditions, where inoculation with some strains reduced the Ni concentration compared to uninoculated plants. Concentrations of all tested elements were mostly in the range of common values for plants. In the soil with a higher content of Ni and Cr, significantly higher concentrations of these elements were found in all treatments, but a greater mass of alfalfa was also measured. The results indicate that inoculation of alfalfa with effective rhizobial strains showed potential for improvement of plants growth and phytostabilisation of trace elements in the soils. Further research is necessary to enable a better explanation of the relationships in the soil-plant-microorganisms system.

KEYWORDS

Soil, rhizobia, trace elements, alfalfa, soil quality.

ACKNOWLEDGEMENTS

This research was supported by the Science Fund of the Republic of Serbia, #GRANT No 7015, Utilizing rhizobia to reduce the risk of heavy metal accumulation in alfalfa: Nickel (Ni) case study – RhizoDETOX and by the Ministry of Science, Technological Development and Innovations of the Republic of Serbia, contract No. 451-03-136/2025-03/200011.