

ISSN 1821-1046

UDK 630

INSTITUTE OF FORESTRY  
BELGRADE




INSTITUT ZA ŠUMARSTVO  
BEOGRAD

**SUSTAINABLE FORESTRY    ODRŽIVO ŠUMARSTVO**

COLLECTION  
TOM 65-66

ZBORNİK RADOVA  
TOM 65-66



**BELGRADE    BEOGRAD**  
**2012.**

ISSN 1821-1046



9 771821 104000

ISSN 1821-1046  
UDK 630

**INSTITUTE OF FORESTRY  
BELGRADE**



**INSTITUT ZA ŠUMARSTVO  
BEOGRAD**

**SUSTAINABLE FORESTRY    ODRŽIVO ŠUMARSTVO**

COLLECTION  
TOM 65-66

ZBORNİK RADOVA  
TOM 65-66

**BELGRADE    BEOGRAD  
2012.**



**INSTITUTE OF FORESTRY BELGRADE** **INSTITUT ZA ŠUMARSTVO BEOGRAD**  
**PROCEEDINGS** **ZBORNİK RADOVA**

**Publisher**

Institute of Forestry  
Belgrade, Serbia

**Izdavač**

Institut za šumarstvo  
Beograd, Srbija

**For Publisher**

Ljubinko Rakonjac, Ph.D.

**Za izdavača**

Dr Ljubinko Rakonjac

**Editorial Board**

Snežana Rajković, Ph.D.  
Institute of Forestry, Belgrade  
Dragana Dražić, Ph.D.  
Institute of Forestry, Belgrade  
Ljubinko Rakonjac, Ph.D.  
Institute of Forestry, Belgrade  
Mara Tabaković-Tošić, Ph.D.  
Institute of Forestry, Belgrade  
Miloš Koprivica, Ph.D.  
Institute of Forestry, Belgrade  
Radovan Nevenić, Ph.D.  
Institute of Forestry, Belgrade  
Mihailo Ratknić, Ph.D.  
Institute of Forestry, Belgrade  
Zoran Miletić, Ph.D.  
Institute of Forestry, Belgrade  
Milorad Veselinović, Ph.D.  
Institute of Forestry, Belgrade  
Biljana Nikolić, Ph.D.  
Institute of Forestry, Belgrade  
Vera Lavadinović, Ph.D.  
Institute of Forestry, Belgrade  
Vesna Golubović-Čurguz, Ph.D.  
Institute of Forestry, Belgrade  
Assoc. Prof. Iantcho Naidenov, Ph.D  
Bulgaria  
Prof. Nikola Hristovski, Ph.D.  
Macedonia  
Dr Kalliopi Radoglou, Ph.D.  
Greece

**Redakcioni odbor**

Dr Snežana Rajković  
Institut za šumarstvo, Beograd  
Dr Dragana Dražić  
Institut za šumarstvo, Beograd  
Dr Ljubinko Rakonjac  
Institut za šumarstvo, Beograd  
Dr Mara Tabaković-Tošić  
Institut za šumarstvo, Beograd  
Dr Miloš Koprivica  
Institut za šumarstvo, Beograd  
Dr Radovan Nevenić  
Institut za šumarstvo, Beograd  
Dr Mihailo Ratknić  
Institut za šumarstvo, Beograd  
Dr Zoran Miletić  
Institut za šumarstvo, Beograd  
Dr Milorad Veselinović  
Institut za šumarstvo, Beograd  
Dr Biljana Nikolić  
Institut za šumarstvo, Beograd  
Dr Vera Lavadinović  
Institut za šumarstvo, Beograd  
Dr Vesna Golubović-Čurguz  
Institut za šumarstvo, Beograd  
Assoc. Prof. Dr Iantcho Naidenov  
Bugarska  
Prof. Dr Nikola Hristovski  
Makedonija  
Dr Kalliopi Radoglou  
Grčka

**Chief Editor**

Snežana Rajković, Ph.D.

**Glavni i odgovorni urednik**

Dr Snežana Rajković

**Technical Editor and Layout**

Tatjana Ćirković-Mitrović, M.Sc.

**Tehnički urednik i prelom teksta**

Mr Tatjana Ćirković-Mitrović

**Secretary**

Tatjana Ćirković-Mitrović, M.Sc.

**Sekretar Zbornika**

Mr Tatjana Ćirković-Mitrović

**Printed in**

100 copies

**Tiraž**

100 primeraka

**Printed by**

Black and White  
Beograd

**Štampa**

Black and White  
Beograd

All rights reserved. No part of this publication might be reproduced by any means: electronic, mechanical, copying or otherwise, without prior written permission of the publisher.

Belgrade, 2012

Preuzimanje članaka ili pojedinih delova ove publikacije u bilo kom obliku nije dozvoljeno bez odobrenja

Beograd, 2012

**Cover Page:** Author of the Photos Tatjana Ćirković-Mitrović, M.Sc.  
**Naslovna strana:** Autor fotografije mr Tatjana Ćirković-Mitrović

CIP - Каталогизација у публикацији  
Народна библиотека Србије, Београд

630

SUSTAINABLE Forestry : collection =  
Održivo šumarstvo = zbornik radova /  
chief

editor = glavni i odgovorni urednik  
Snežana

Rajković. - 2008, T. 57/58- . - Belgrade  
(Kneza Višeslava 3) : Institute of forestry,  
2008- (Beograd : Black and White). - 24  
cm

Godišnje. - Je nastavak: Zbornik radova -  
Institut za šumarstvo = ISSN 0354-1894  
ISSN 1821-1046 = Sustainable Forestry  
COBISS.SR-ID 157148172

# CONTENT SADRŽAJ

## TOM 65-66

---

Vera LAVADINOVIĆ, Nenad MARKOVIĆ

**THE DIFFERENCES OF NEEDLE LENGTH OF DOUGLAS-FIR  
PROVENANCES AT TWO SITES IN TEST PLANTATIONS**

7

---

Vladan POPOVIĆ, Mirjana ŠIJAIĆ-NIKOLIĆ, Ljubinko RAKONJAC,  
Dušan JOKANOVIĆ

**VARIABILITY IN CONE MORPHOMETRIC CHARACTERS  
AMONG TEST TREES OF BALD CYPRESS (*Taxodium distichum* L.  
Rich.) IN SEED STAND NEAR BAČKA PALANKA**

15

---

Vlado ČOKEŠA, Miloš KOPRIVICA, Snežana STAJIĆ, Zoran MILETIĆ,  
Bratislav MATOVIĆ

**NATURAL REGENERATION OF BEECH FORESTS IN SERBIA**

27

---

Mihailo RATKNIĆ, Ljubinko RAKONJAC, Tatjana RATKNIĆ,  
Sonja BRAUNOVIĆ

**BEECH AND FIR FOREST RESOURCES IN THE PEŠTER PLAIN  
AREA**

41

---

Svetlana BILIBAJKIĆ, Tomislav STEFANOVIĆ, Radovan NEVENIĆ,  
Zoran PODUŠKA, Renata GAGIĆ SERDAR, Ilija ĐORĐEVIĆ,  
Goran ČEŠLJAR, Zoran MILETIĆ

**INTENSIVE MONITORING AT LEVEL II TEST PLOT KOPAONIK  
IN 2012**

51

---

Sonja BRAUNOVIĆ, Mihailo RATKNIĆ, Ljubinko RAKONJAC

**IMPACT OF THE PARENT ROCK ON EROSION PROCESS  
DEVELOPMENT IN GRDELICA GORGE AND VRANJE VALLEY**

65

---

Tatjana ĆIRKOVIĆ-MITROVIĆ, Vladan POPOVIĆ,  
Ljiljana BRAŠANAC-BOSANAC, Ljubinko RAKONJAC,  
Aleksandar LUČIĆ

**THE EFFECT OF APPLICATION OF DIFFERENT NUTRITION  
PREPARATIONS ON HEIGHT INCREMENT OF WALNUT  
(*Juglans regia* L.) SEEDLINGS**

73

---

Milorad VESELINOVIĆ, Vesna GOLUBOVIĆ-ĆURGUZ,  
Suzana MITROVIĆ, Dragana DRAŽIĆ, Nevena ČULE, Biljana NIKOLIĆ,  
Đorđe JOVIĆ

**MICROBIOLOGICAL ACTIVITY OF THE FOREST SOIL IN THE  
AREAS WITHIN THE TERRITORY OF BELGRADE**

83

---

Miroslava MARKOVIĆ, Snezana RAJKOVIĆ, Radoslav RAJKOVIĆ <b>MANALYSIS OF THE IMPACT OF INJURIES CAUSED BY THE INFLUENCE OF MECHANICAL AND ABIOTIC FACTORS ON THE OCCURRENCE OF HARMFUL FUNGAL ORGANISMS</b>	97
Vesna GOLUBOVIĆ ĆURGUZ, Katarina MLADENOVIĆ, Biljana NIKOLIĆ, Snežana RAJKOVIĆ <b>DISEASES AND INJURIES OF HIMALAYAN PINE SEED FROM CULTURE GROWN ON DEPOSOL</b>	111
Slobodan MILANOVIĆ, Zlatan RADULOVIĆ, Milorad VESELINOVIĆ, Suzana MITROVIĆ, Katarina MLADENOVIĆ <b>SENSITIVITY OF SEVEN CLONES OF POPLAR TO THE ATTACK BY CATERPILLARS OF GYPSY MOTH (<i>Lymantria dispar</i> L.) AND FUNGUS <i>Pollacia elegans</i> (Vuill.) Fabr.</b>	123
Mara TABAKOVIĆ-TOŠIĆ <b>GYPSY MOTH, <i>Lymantria dispar</i> (L.), AND ITS NATURAL ENEMIES IN THE FORESTS OF CENTRAL SERBIA</b>	133
Renata GAGIĆ SERDAR, Radovan NEVENIĆ, Svetlana BILIBAJKIĆ, Tomislav STEFANOVIĆ, Zoran PODUŠKA, Ilija ĐORĐEVIĆ, Goran ČEŠLJAR <b>THE MOST INVASIVE FIVE WOODY INVASIVE PLANTS CONTROL RISKS ASSESSMENT - STRATEGIC PRECONDITION FOR SUSTAINABLE GOVERNANCE OF NATURAL RESOURCES IN SERBIA</b>	149

UDK 630\*416.5+630\*443.3=111  
Original scientific paper

**MANALYSIS OF THE IMPACT OF INJURIES CAUSED  
BY THE INFLUENCE OF MECHANICAL AND ABIOTIC FACTORS  
ON THE OCCURRENCE OF HARMFUL FUNGAL ORGANISMS**

Miroslava MARKOVIC<sup>1</sup>, Snezana RAJKOVIC<sup>1</sup>, Radoslav RAJKOVIC<sup>2</sup>

***Abstract:** The research described in this paper is focused on the occurrence of pathogenic microorganisms on beech trees relative to the presence of tree injuries, with the aim to ensure protection and preservation of this species in Serbia. The research was conducted in eastern Serbia, in a hillside beech forest *Fagetum moesiacaе submontanum* of generative origin. The testing was carried out on two sites over 51 testing plots, with a total of 829 trees and 21 species of identified fungi. On the first site it was found that the appearance of fungi primarily depends on the presence of mechanical damage on trees (as much as 73.46%), while the presence of abiotic damage has almost no bearing (only 3.21%). On the second site there was a strong correlation link between the occurrence of fungi and presence of mechanical damage - 51.88%, as well as between the fungi and abiotic damage – 47.96%. The health condition of high beech stands was found to be heavily dependent on careful and proper manipulation during harvesting, while each injury inflicted on a beech live tree during logging opens the door to infection with pathogenic microorganisms.*

---

<sup>1</sup>PhD Miroslava Markovic, Research Associate, Institute for Forestry, Kneza Visislava 3, 11030 Belgrade, Serbia, Mob. tel.: +381691999116, Fax.: +381112545969, Email: mira013@gmail.com

<sup>1</sup>PhD Snezana Rajkovic, Principal Research Fellow, Institute for Forestry, Kneza Visislava 3, 11030 Belgrade, Serbia

<sup>2</sup>MSc Radoslav Rajkovic, Research Assistant, University of Belgrade, Faculty of Mechanical Engineering, Queen Mary 16, 11030 Belgrade, Serbia

\* **Acknowledgement:** *The study was carried out within the Project TP-31070: “The development of technological methods in forestry in order to attain optimal forest cover”, financed by the Ministry of Education and Science of the Republic of Serbia within the framework of integrated and interdisciplinary research for the period 2011 – 2014.*

**Key words:** injuries, environmental factors, fungi, beech forests

## АНАЛИЗА УТИЦАЈА ОЗЛЕДА ПРОУЗРОКОВАНИХ ДЕЈСТВОМ МЕХАНИЧКИХ И АБИОТИЧКИХ ФАКТОРА НА ПОЈАВУ ШТЕТНИХ ГЉИВИЧНИХ ОРГАНИЗАМА

**Извод:** У раду је истраживан аспект појаве патогених микроорганизама на буковом дрвету у односу на присуство озледа на стаблима, у циљу заштите и очувања ове врсте у Србији. Истраживања су вршена у источној Србији, у брдској шуми букве *Fagetum moesiacaе submontanum*, генеративног порекла. Испитивањем је обухваћено два локалитета на 51 огледној парцели, са укупно 829 стабала и констатовано је присуство 21 врста гљива. На првом локалитету констатовано је да појава гљива првенствено зависи од присуства механичких оштећења на стаблима (чак 73.46%), а готово уопште није у вези са присуством абиотичких оштећења (свега 3.21%). На другом локалитету постоји јака корелациона веза између појаве гљива и присуства механичких оштећења – 51.88%, као и између гљива и абиотичких оштећења – 47.96%. Констатовано је да за здравствено стање високих букових састојина изузетан значај има пажљиво и правилно манипулисање приликом сече, а свака озледа на буковим дубећим стаблима почињена при сечи је отворен пут за заразу патогеним микроорганизмима.

**Кључне речи:** оштећења, спољна средина, гљиве, букове шуме

### 1. INTRODUCTION

The strategy of preserving the biodiversity and genetic resources of economically most valuable species of trees dictates that Serbia's forestry should primarily be concerned with preservation of the abundance of natural forests as a national wealth (Milovanovic *et al* 2004). According to the internationally adopted definition, sustainable forest management means “*the stewardship and use of forests and forest lands in a way, and at a rate, that maintains their biodiversity, productivity, regeneration capacity, vitality and their potential to fulfil relevant ecological, economic and social functions, and that does not cause damage to other ecosystems*” (MCPFE, Helsinki 1993, as cited in Medarevic *et al* 2008).

Serbia's forest area relative to the global aspect is close to the world's level of 30%, but significantly below the European average of 46%. Out of the total 29.1% of forest area in Serbia, 7.1% is in Vojvodina whereas 37.6% of forests are located in Central Serbia (Bankovic *et al* 2009). The national inventory of Serbia's forests in the total volume and bulk growth is dominated by beech, whose presence amounts to 42.4%, or 32.3% (Bankovic *et al* 2000).

Due to their presence in the forest reserves of Serbia, beech forests undoubtedly have the greatest significance. We can therefore conclude, quite justifiably, that management of beech forests is a much more complex and difficult task compared to management of any other tree species. In addition, available references most frequently speak of the quality of tall beech forests in descriptive and general terms – that it is unsatisfactory and in need of improvement (Stojanovic & Krstic 2003, Koprivica *et al* 2009).

The biological properties, ecological demands, natural distribution, stewardship values and generally beneficial functions of beech forests, along with their structure, make beech the basic tree species for Serbian forestry (Vuckovic *et al* 2005), although the use of beech lumber on a wider scale is limited by its short lifespan.

Beech wood is vulnerable and represents an excellent base for development of numerous parasitic and saprophytic organisms, among which primarily parasitic fungi and harmful insects. In beech coppice forests in Serbia, the total of 147 species of fungi were found on beech trees, out of which 33 species occur on crowns, fruits and young crop, 56 species occur on leaves and bark of the branches and the trunk, whereas 58 species of fungi cause rot and coloration of wood (Karadzic & Milijasevic 2005).

The cause of beech forests dieback is a consequence of simultaneous negative impact of climatic (climate changes), management and biotic factors. Among these a special place belongs to man, whose irrational exploitation of beech forests resulted in Serbia's area under forest being cut almost in half. Deforestations of beech woodland that occurred in the past (in particular immediately after World War II) were not at all conducted as regeneration harvests, but almost exclusively for exploitation purposes. As a consequence of such management practices, forests have become extremely sensitive to harmful effects of numerous abiotic and biotic factors, notably parasitic fungi and harmful insects among the latter. The problem of protection of beech forests is further complicated by the occurrence of dangerous diseases and a large number of wood destructors that start their development as parasites on living trees and continue as saprophytes on timber (Tabakovic-Tosic & Markovic 2003, Miletic *et al* 2006, Markovic *et al* 2011a, Markovic *et al* 2011b).

This paper researched one aspect of occurrence of pathogenic microorganisms on beech trees, with the aim to contribute to the most rational approach to use of beech timber while preserving the beech stands in Serbia to the maximum extent.

## **2. MATERIAL AND METHODS**

The sites selected for research were the ones on which the observation method revealed a large number of injuries on trees. The paper provides an analysis of the impact of tree injuries on occurrence of pathogenic and epixylic fungi on live trees in beech woodlands.

The research was carried out in the forest holding "Severni Kucaj" in Kucevo, forest administration Kucevo, Eastern Serbia, in a hillside forest of *Fagetum moesiaca submontanum* beech of generative origin. The first tested site was located in the administration unit Majdan Kucajna, division 33. The second site was in administration unit Crni Vrh, division 42, sections *a* and *b*. The research included the total of 829 beech trees on 51 test plots.

The 500 m<sup>2</sup> trial experimental plots were circular, placed in the stands at 100 x 100 m distances (according to the method described by Koprivica *et al* 2008). Each experimental plot included between 4 and 24 trees on site I, and

between 8 and 27 trees on site II. Injuries noted on each tree were classified as mechanic (injuries from felling and hauling during harvest) and abiotic (injuries from wind, snow, ice, frost and excessive insulation that caused bark inflammation). The methods used were those described by Koprivica & Matovic, 2005, Markovic *et al* 2007. In addition, any presence of pathogenic and epyxilic fungi on trees was also noted. On the basis of the received data, statistical analysis was conducted in order to determine the correlation link.

### 3. RESULTS AND DISCUSSION

Table 1 presents an overview of the fungi identified on site 1 according to their frequency of occurrence. It is evident from the table that the first 4 fungi are present in all experimental plots (noted on 12.6% to 29% of all trees), whereas the presence of the latter 8 fungi was noted, on the average, in only 4 plots and no more than 0.3% to 0.9% of trees. The fungi with ordinal numbers 5 through 9 are present on over 50% of experimental plots (or 6.5% to 11.1% of trees), while the fungi with ordinal numbers 8 through 13 were identified on less than 50% of the plots and spread on 1% to 20% of the tested trees.

Table 1 and table 2 presents fungi classified according to their significance, where those with 3 stars represent dangerous fungi with high significance, the fungi with 2 stars have medium significance, the fungi with one star have low significance, and those without stars have no significance.

**Table 1.** *Fungi identified on site I – Administration unit Majdan Kucajna, division 33*

Ordinal number of fungus	Type of fungus	Significance of fungus	% plots on which the fungus is present	% trees on which the fungus is present
1	<i>Apiognomonía errabunda</i>	***	100.0	29.0
2	<i>Coriolus sp.</i>	**	100.0	24.6
3	<i>Hypoxylon sp.</i>	**	100.0	8.9
4	<i>Stereum sp.</i>	**	100.0	12.6
5	<i>Diatrype stigma</i>	*	65.2	11.1
6	<i>Fomes fomentarius</i>	***	57.0	6.5
7	<i>Trametes sp.</i>	**	52.2	8.6
8	<i>Diatrype disciformis</i>	**	47.8	20.9
9	<i>Nectria galligena</i>	***	30.4	4.0
10	<i>Armillaria mellea</i>	***	26.0	1.2
11	<i>Lenzites trabaeae</i>	**	17.4	1.8
12	<i>Nectria coccinea</i>	***	8.7	1.2
13	<i>Fomes igniarius</i>	**	8.7	1.5
14	<i>Pleurotus ostreatus</i>	***	4.3	0.6
15	<i>Poria obliqua</i>	***	4.3	0.3

16	<i>Dedalea quercina</i>	**	4.3	0.9
17	<i>Exidia recisa</i>	*	4.3	0.3
18	<i>Hydnum coraloides</i>	*	4.3	0.3
19	<i>Auricullaria auricula judae</i>	-	4.3	0.3
20	<i>Auricullaria mesenterica</i>	-	4.3	0.3
21	<i>Bulgaria polymorpha</i>	-	4.3	0.6

Under the classification proposed by Karadzic 2003, the present fungi were classified as follows:

\*\*\* fungi with high significance (the highest significance is given to fungi that act that both as parasites and saprophytes, i.e. whose activity starts on standing, live trees and then persists on dead trees, following the harvest). These species of epyxilic fungi demonstrate a very high level of destruction and degrade primarily lignin, as well as cellulose and hemicelluloses, but to a lesser degree. Among the identified fungi, this group comprises *Armillaria mellea*, *Fomes fomentarius*, *Pleurotus ostreatus* and *Poria obliqua*. Besides the above-named wood-decay fungi, this group also includes the following pathogenic fungi: *Apiognomonium errabunda*, *Nectria coccinea* and *Nectria galligena*).

\*\* fungi with medium significance (this group comprises the fungi that cause a somewhat lower degree of destruction, but appear on both injured, weakened trees and the freshly harvested ones). This group is represented by *Coriolus versicolor*, *Dedalea quercina*, *Diatrype disciformis*, *Hypoxylon* sp., *Stereum* sp. and *Trametes* sp.

\* fungi with low significance (this group comprises the fungi that appear on rotting trees, frequently causing their complete degradation). Among the identified fungi, this group includes *Exidia recisa*, *Hydnum* sp. and *Diatrype stigma*.

- fungi with no significance (representatives of this group identified on the tested sites include *Auricullaria auricula judae*, *Auricullaria mesenterica* and *Bulgaria polymorpha*).

**Table 2.** Fungi identified on site II – Administration unit Crni Vrh, division 42, sections a, b

Ordinal number of fungus	Type of fungus	Significance of fungus	% plots on which the fungus is present	% trees on which the fungus is present
1	<i>Coriolus</i> sp.	**	100.0	18.3
2	<i>Apiognomonium errabunda</i>	***	82.0	15.0
3	<i>Hypoxylon</i> sp.	**	78.6	6.5
4	<i>Nectria coccinea</i>	***	64.3	7.1
5	<i>Nectria galligena</i>	***	53.6	6.5
6	<i>Diatrype stigma</i>	*	35.7	5.7

7	<i>Fomes fomentarius</i>	***	21.0	2.2
8	<i>Hydnum coraloides</i>	*	14.3	0.8
9	<i>Stereum sp.</i>	**	10.7	1.2
10	<i>Diatrype disciformis</i>	**	7.1	1.4
11	<i>Armillaria mellea</i>	***	7.0	0.6
12	<i>Dedalea quercina</i>	**	3.6	0.2

Table 3 presents the testing results for the total number of the present fungi and the total number of mechanical and abiotic damage on site II.

One of the most significant fungi identified on the tested sites is *Nectria coccinea* (Pers. Ex Fr.) Fries., which together with the insect *Cryptococcus fagisuga* Lind. causes the so-called “beech bark disease”. On site I, this fungus was found on 1.2% of trees or on 3 experimental plots (1,3 and 9), whereas on site II its spread was much greater and covered 7.1% of trees or almost two-thirds of experimental plots. This disease is lately being regarded as a major factor compromising normal development of beech trees, which merits special attention given the fact that it is spreading over ever-larger areas.

**Table 3.** Overview of attack by fungi and injuries on beech trees on site I – Administration unit Majdan Kucajna, division 33

(x)	Number of fungi found			(x <sub>1</sub> )	(x <sub>2</sub> )	(x <sub>3</sub> )	Index (x <sub>3</sub> /x)
	(y <sub>1</sub> )	(y <sub>2</sub> )	(y <sub>3</sub> )				
No. of trees on the plot	No. of dangerous fungi	No. of other fungi	Total number of fungi	Number of mechanical injuries (damage from hauling and felling during harvest)	Number of abiotic injuries (damage from wind, snow, ice, frost and excessive insulation – bark inflammation)	Total number of mechanical and abiotic injuries	
12	1	3	4	2	3	5	0.42
24	1	4	5	3	4	7	0.29
9	1	3	4	2	3	5	0.56
11	1	3	4	1	3	4	0.36
16	1	5	6	8	1	9	0.56
9	1	4	5	6	2	8	0.89
16	4	6	10	15	6	21	0.06
12	4	6	10	20	9	29	2.42
23	6	9	15	27	4	31	1.35
15	1	6	7	3	1	4	0.27
21	2	6	8	17	9	26	1.24
18	3	6	9	21	3	24	1.33
8	1	5	6	14	2	16	2.00
11	2	6	8	15	3	18	1.64
19	2	6	8	11	10	21	1.11
4	1	5	6	4	1	5	1.25
15	3	5	8	16	42	20	1.33
9	1	4	5	9	6	11	1.22
23	4	8	12	19	7	25	1.09

8	2	4	6	7	3	14	1.75
8	1	5	6	14	3	17	2.12
17	4	6	10	12	8	20	1.18
16	3	7	10	23	4	27	1.69

Measures undertaken against this fungus are classified into several categories:

- biological preventive measures, including use of predators and super-parasites against insects (prior to infection with fungus),
- bio-control of the fungus by means of antagonists (once the infection occurs),
- silvicultural measures – removal of diseased trees (in advanced stages of the infection),
- chemical measures, which are non-economical for forests and thus applied only to parks and alleys of trees.

It is important to note that following the infection of beech trees with this fungus, the necrotic bark sections very quickly get infested by wood-decaying fungi and wood-destroying insects, which also play a role in rapid tree decay and extinction of beech trees (Karadzic 2003, Ivkovic et al 2007).

The data presented in Table 3 served as basis for performance of a statistical analysis – simple and multiple linear regression between all pairs in the presented columns, and correlation matrixes made between columns  $x$ ,  $y_1$ ,  $y_2$  and  $y_3$ , as well as columns  $x_1$ ,  $x_2$ ,  $x_3$  and  $x_3/x'$ . The correlation analysis clearly demonstrates that in all cases there is a link between the number of trees ( $x$ ) and other columns. Next, there is a correlation link between the number of dangerous fungi ( $y_1$ ) and other columns, with the exception of abiotic injuries ( $x_2$ ) and index representing a quotient between the total number of injuries and the number of trees ( $x_3/x'$ ). The same applies to columns  $y_2$  (other fungi),  $y_3$  (total number of fungi) and  $x_1$  (mechanical injuries). Column  $x_2$  has no correlation links with any other column, whereas  $x_3$  (the total number of mechanical and abiotic injuries) has links to all columns except abiotic injuries ( $x_2$ ). Column  $x_3/x'$  (index) is not linked to other columns, except to columns  $x_1$  and  $x_3$  (mechanical injuries and total number of injuries).

This practically means that, on site I, the occurrence of fungi (both dangerous and other) – column  $y_3$  is primarily contingent upon the presence of mechanical injuries -  $x_1$  (as much as 73.46%), while the remaining 26.54% depends on other factors – tree condition (susceptibility to disease), position inside the stand (open trees or within a dense canopy, land elevation, geological base, etc.), climatic conditions during the year that may or may not favour the development of fungi, etc. On the other hand, statistical analysis of the data received from site I shows that the occurrence of fungi is not linked to damage caused by activity of abiotic factors (the correlation link is very low at 3.21%).

Therefore, careful and proper handling of trees during felling is critical for the health condition of tall beech stands. Every injury sustained by live beech trees during felling opens the door to infection by pathogenic microorganisms.

Condition on site II is presented in Table 4, which was the basis for performance of a statistical analysis – simple and multiple linear regression

between all pairs in the presented columns and correlation matrixes made between columns  $x$ ,  $y_1$ ,  $y_2$  and  $y_3$ , as well as columns  $x_1$ ,  $x_2$ ,  $x_3$  and  $x_3/x$ .

Statistical analysis of the obtained data demonstrated that on this site there was a significant statistical link between all presented columns, as well as between the number of abiotic injuries and occurrence of fungi. While this was not the case on the previous site, it appeared here as a consequence of a large number of injuries. The strongest correlation link was the one between the total number of fungi and mechanical injuries (columns  $y_3$  and  $x_1$ ), amounting to 51.88%. Another strong link existed between the total number of fungi and abiotic injuries (columns  $y_3$  and  $x_2$ ), only slightly weaker than the previous one at 47.96%. The links between the number of dangerous and other fungi ( $y_1$ ,  $y_2$ ) and mechanical and abiotic injuries ( $x_1$ ,  $x_2$ ) were significant, ranging from 26.70% and 36.47%, where the links between the fungi and mechanical injuries were stronger by roughly 2 to 5% than the links between the occurrence of fungi and abiotic injuries.

**Table 4.** Overview of attack by fungi and injuries on beech trees on site II – Administration unit Crni Vrh, division 42, sections a, b

$(x)$ No. of trees on the plot	Number of fungi found			$(x_1)$ Number of mechanical injuries (damage from hauling and felling during harvest )	$(x_2)$ Number of abiotic injuries (damage from wind, snow, ice, frost and excessive insulation – bark inflammation)	$(x_3)$ Total number of mechanical and abiotic injuries	Index ( $x_3/x$ )
	$(y_1)$ No. of dangerous fungi	$(y_2)$ No. of other fungi	$(y_3)$ Total number of fungi				
20	5	6	11	16	42	58	2.90
27	3	3	6	12	7	19	0.70
22	5	3	8	16	34	50	2.27
20	4	2	6	4	34	38	1.90
21	3	3	6	10	17	27	1.29
9	1	2	3	4	8	12	1.33
11	0	3	3	7	7	14	1.27
14	0	2	2	2	7	9	0.64
22	3	2	5	5	7	12	0.54
17	4	2	6	8	37	86	5.06
10	1	2	3	2	19	19	1.90
12	2	2	4	8	14	22	1.83
16	2	2	4	10	7	17	1.06
23	3	2	5	10	3	13	0.56
21	3	2	5	14	2	16	0.76
25	2	5	7	16	27	43	1.72
19	1	3	4	11	9	20	1.05
14	1	2	3	3	11	14	1.00
23	4	2	6	14	10	24	1.04
19	3	2	5	5	17	22	1.16
11	0	2	2	6	8	14	1.27
23	3	2	5	1	18	19	0.83
20	1	2	3	3	9	12	0.60
20	0	2	2	1	3	4	0.20
8	0	2	2	1	15	16	2.00
26	3	2	5	12	20	32	1.23
18	3	2	5	15	15	30	1.67
13	3	2	5	7	8	15	1.15

This means that the occurrence of fungi (both dangerous and other fungi) on the second site – column  $y_3$  – was directly linked to the presence of mechanical and abiotic injuries -  $x_1$  and  $x_2$ . In other words, the results obtained through comparative analysis of sites I and II lead to conclusion that the number of injuries is in fact the determining factor linking the occurrence of fungi and damage to trees. On sites with fewer injuries the correlation links between the occurrence of fungi and the injuries on trees are less strong, and vice versa.

It is a well-known fact that health status of the stands is contingent upon a large number of factors, among which year-round climatic conditions must be considered as one of the most critical. Rainy, humid and relatively warm weather favours the activity of the fungi and increases the yield, and thus enables not only faster colonization by the fungi but also more precise identification of the existing microflora. It should also be noted that diagnosis of the disease is greatly impeded by long incubation of the fungi colonizing vital trees, while primary symptoms appear on the surface only after several years of attack (reproductive organs – visible carpophores may not appear at all or their appearance might be extended over a number of years). In addition to an accurate diagnosis, it is essential to make a precise prognosis of the dynamics of development of pathological processes in the plant. However, this prognosis cannot be determined with any reliable level of accuracy for the upcoming calendar years, as climatic conditions are a determining factor for the development of the infection. It is thus possible to make only a rough prognosis, based on mapping the parts of the forest under attack according to the destructor species and attack intensity, and use it as basis for planning the sanitary and silvicultural activities.

Sanitation felling and other phytosanitary measures, which may or may not be carried out in forests, certainly have a great impact on general health condition of the stands. Proper stewardship can minimize the existing infections and thus eliminate or greatly mitigate any new infection, which significantly contributes to having the health status of the stands restored and maintained on a satisfactory level.

#### **4. CONCLUSIONS**

On the first tested site, the occurrence of fungi primarily depended on the presence of mechanical damage, where this link was quite strong with as much as 73.46%, while the remaining 26.54% were contingent upon other factors. On the other hand, statistical analysis of the data received on the first site showed that the occurrence of fungi had almost no connection to the presence of abiotic damage (the correlation link was only 3.21%).

On the second site, there was a statistically significant difference between all columns, as well as between the number of abiotic injuries and the occurrence of fungi. This was not the case on the previous site, but appeared here as a result of a large of number of injuries. The strongest correlation link was the one between the total number of fungi and the mechanical injuries, amounting to 51.88%. The link between the total number of fungi and abiotic injuries was also strong at 47.96%, whereas the links between the number of dangerous and other fungi on one hand and mechanical and abiotic injuries on the other, ranging 26.70% to

36.47%, may be considered significant. This effectively means that the occurrence of fungi on the second site was directly linked to the presence of both mechanical and abiotic injuries.

Based on the results of the comparative analysis of the two sites, the number of injuries may be identified as the determining factor linking the occurrence of fungi and the damage on trees. On sites with fewer injuries the correlation links between the occurrence of fungi and the injuries are less strong, and vice versa.

Careful and proper handling of trees during felling is critical for the health condition of tall beech stands. Every injury sustained by live beech trees during felling opens the door to infection by pathogenic microorganisms. Proper stewardship may minimize the existing infections and thus eliminate or greatly mitigate any new infection, which significantly contributes to having the health status of the stands restored and maintained on a satisfactory level.

## REFERENCES

- Bankovic, S., Medarevic, M., Pantic, D., Petrovic, N., Sljukic, B., Obradovic, S. (2000): *Growing stock of Republic of Serbia – The situation and problems. Bulletin of Faculty of Forestry No. 100, pp. 7-30*
- Bankovic, S., Medarevic, M., Pantic, D., Petrovic, N. (2009): *National Forest Inventory of the Republic of Serbia – Growing stock of Republic of Serbia*. Ministry of Agriculture Forestry and Water Management Serbia - Management of Forests, Belgrade, Serbia, pp. 1-244
- Ivkovic, S., Markovic, M., Miric, M (2007): *Growth and mycelial mass production of wood – decaying fungus *Trametes versicolor* (Fr.) Pil. and *Coniophora puteana* (Schum.:Fr.) Karst in controlled temperature and pH conditions*. International Scientific Conference "Integral protection of Forests Scientific – Technological platform", Belgrade, 12.12.2007, Book of abstracts, pp 95-96
- Karadzic, D. (2003): *The most important disease of the beech forests in Serbia. Forestry No. 1-2, pp. 59-72*
- Karadzic, D, Milijasevic, T. (2005): *Beech (*Fagus moesiaca* /Domin, Mally/ Czezzott.) in Serbia*. Monography, Chapter 3, Faculty of Forestry, Belgrade, Serbia, pp 179-196
- Koprivica, M., Matovic, B. (2005): *Regression equations of volume and volume increment of beech trees in high forests in Serbia. Sustainable Forestry, No. 52-53, pp. 5-17* [in Serbian]
- Koprivica, M., Matovic, B., Markovic, N. (2008): *Qualitative and assortment structure of high beech stand volume in Severnokucajsko forest area. Forestry, Vol. 60, No. 1-2, pp. 41-52* [in Serbian]
- Koprivica, M., Matovic, B., Cokesa, V. (2009): *Qualitative structure of volume, volume increment and yield of a beech high stand. Forestry, Vol. 61, Iss. 1-2, pp 95-108*
- Markovic, N., Koprivica, M., Matovic, B. (2007): *Software "Assortment" designed for processing field data*. Institute for Forestry, Belgrade, Serbia [in Serbian]

Markovic, M., Rajkovic, S., Miric, M., Mitic, D., Milovanovic, J., Tabakovic-Tosic, M. (2011a): *Colonization of the substrate of wood – decaying fungi Fomitopsis pinicola (Sw.:Fr.) P. Karst. isolated from beech and fir under controlled temperature and pH conditions. Fresenius Environmental Bulletin, Vol. 20, Issue 3, pp 583-589*

Markovic, M., Rajkovic, S., Miric, M., Mitic, D., Rakonjac, Lj. (2011b): *Growth conditions of mycelium medicinal mushroom Lentinula edodes (Berk.) Pegl. in the substrate colonization phase”, Scientific Research and Essays, 8 September, Vol 6(19), pp 4133-4140*

Medarevic, M., Bankovic, S., Sljukic, B. (2008): *Sustainable Forest Management in Serbia - State and Potentials. Bulletin of the Faculty of Forestry 97, pp 33-56*

Miletic, Z., Markovic, M., Tabakovic-Tosic, M. (2006): *Effect of Different Soil Types on Hazard to Beech Caused by the Fungi in the Genus Nectria in M.U. "Majdan-Kucajna". International Scientific Conference in occasion of 60 year of operation of Institute of Forestry, Belgrade, Serbia. Sustainable Use of Forest Ecosystems-The Challenge of the 21<sup>st</sup> Century, 8-10 November, Donji Milanovac, Proceedings, pp. 297-302*

Milovanovic, D., Krstic, M., Babic, V. (2004): *Significance of productive and social functions for the beech forests in Serbia. Forestry, No 1-2, January-June, pp 105-116*

Stojanovic, Lj., Krstic, M. (2003): *Basic problems of growing the beech forests. Forestry, vol. 55, No. 1-2, pp 25-38*

Tabakovic-Tosic, M., Markovic, M. (2003): *Harmful biotic factors in beech copice forests of east Serbia. International scientific conference " 75 years of the Forest Research Institute of Bulgarian Academy of Sciences" , 1-5 October. Sofia. Bulgaria Proceedings of scientific papers 2, pp. 239-246*

Vuckovic, M., Stajic, B., Radakovic, N. (2005): *Significance of the monitoring diameter increment in terms of bioindication vitality of stems and stands. Forestry, Vol. 57, No. 1-2, pp 1-10*

## **ANALYSIS OF THE IMPACT OF INJURIES CAUSED BY THE INFLUENCE OF MECHANICAL AND ABIOTIC FACTORS ON THE OCCURRENCE OF HARMFUL FUNGAL ORGANISMS**

*Miroslava MARKOVIC, Snezana RAJKOVIC, Radoslav RAJKOVIC*

### **Summary**

The research was conducted in eastern Serbia, in a hillside beech forest *Fagetum moesiaca submontanum* of generative origin. The testing was carried out on two sites over 51 testing plots, with a total of 829 trees and 21 species of identified fungi. On the first site it was found that the appearance of fungi primarily depends on the presence of mechanical damage on trees (as much as 73.46%), while the presence of abiotic damage has almost no bearing (only 3.21%). On the second site there was a strong correlation link between the occurrence of fungi and presence of mechanical damage - 51.88%, as well as between the fungi and abiotic damage – 47.96%.

On the first tested site, the occurrence of fungi primarily depended on the presence of mechanical damage, where this link was quite strong with as much as 73.46%, while the remaining 26.54% were contingent upon other factors. On the other hand, statistical

analysis of the data received on the first site showed that the occurrence of fungi had almost no connection to the presence of abiotic damage (the correlation link was only 3.21%).

On the second site, there was a statistically significant difference between all columns, as well as between the number of abiotic injuries and the occurrence of fungi. This was not the case on the previous site, but appeared here as a result of a large of number of injuries. The strongest correlation link was the one between the total number of fungi and the mechanical injuries, amounting to 51.88%. The link between the total number of fungi and abiotic injuries was also strong at 47.96%. This effectively means that the occurrence of fungi on the second site was directly linked to the presence of both mechanical and abiotic injuries.

Based on the results of the comparative analysis of the two sites, the number of injuries may be identified as the determining factor linking the occurrence of fungi and the damage on trees. On sites with fewer injuries the correlation links between the occurrence of fungi and the injuries are less strong, and vice versa. The health condition of high beech stands was found to be heavily dependent on careful and proper manipulation during harvesting, while each injury inflicted on a beech live tree during logging opens the door to infection with pathogenic microorganisms.

## АНАЛИЗА УТИЦАЈА ОЗЛЕДА ПРОУЗРОКОВАНИХ ДЕЈСТВОМ МЕХАНИЧКИХ И АБИОТИЧКИХ ФАКТОРА НА ПОЈАВУ ШТЕТНИХ ГЉИВИЧНИХ ОРГАНИЗАМА

*Мирослава МАРКОВИЋ, Снежана РАЈКОВИЋ, Радослав РАЈКОВИЋ*

### Резиме

Истраживања су вршена у источној Србији, у брдској шуми букве *Fagetum moesiacaе submontanum*, генеративног порекла. Испитивањем је обухваћено два локалитета на 51 огледној парцели, са укупно 829 стабала и констатовано је присуство 21 врста гљива. На првом локалитету констатовано је да појава гљива првенствено зависи од присуства механичких оштећења на стаблима (чак 73,46%), а готово уопште није у вези са присуством абнотичких оштећења (свега 3,21%). На другом локалитету постоји јака корелациона веза између појаве гљива и присуства механичких оштећења – 51,88%, као и између гљива и абнотичких оштећења – 47,96%.

На првом испитиваном локалитету, појава гљива првенствено зависи од присуства механичких оштећења – веза је јака, износи чак 73,46%, а осталих 26,54% зависи од других фактора. Насупрот томе, статистичка анализа добијених података на првом локалитету показује да појава гљива готово да уопште није у вези са присуством абнотичких оштећења (корелациона веза износи свега 3,21%).

На другом локалитету постоји значајна статистичка веза између свих колона, па и између броја абнотичких оштећења и појаве гљива, што није био случај са претходно приказаним локалитетом, а што је у овом случају последица великог броја оштећења. Најјача корелациона веза постоји између укупног броја гљива и механичких оштећења и износи 51,88%. Веза између укупног броја гљива и абнотичких оштећења је такође јака и износи 47,96%. То практично значи да је појава гљива на другом локалитету, у директној вези са присуством и механичких и абнотичких оштећења.

Ако се посматрају резултати добијени упоредном анализом првог и другог локалитета, може се рећи да је опредељујући фактор који доводи у везу појаву гљива

и оштећења на стаблима управо број оштећења. На локалитетима са мањим бројем оштећења корелационе везе између појаве гљива и озледа на стаблима су слабије и обрнуто. Констатовано је да за здравствено стање високих букових састојина изузетан значај има пажљиво и правилно манипулисање приликом сече, а свака озледа на буковим дубећим стаблима почињена при сечи је отворен пут за заразу патогеним микроорганизмима.

