



2022

ABSTRACT BOOK

INTERNATIONAL SOIL SCIENCE SYMPOSIUM on SOIL SCIENCE & PLANT NUTRITION

(7th International Scientific Meeting)

2 – 3 December 2022

Samsun, Türkiye

Editors

Dr. Rıdvan KIZILKAYA

Dr. Coşkun GÜLSER

Dr. Orhan DENGİZ

Organized by

Federation of Eurasian Soil Science Societies

Erasmus Mundus Joint Master Degree in Soil Science (emiSS) Programme



With the support of the
Erasmus+ Programme
of the European Union

Cover design by FESSS

Editors:

Dr.Rıdvan Kızılkaya

Ondokuz Mayıs University, Faculty of Agriculture
Department of Soil Science and Plant Nutrition
55139 Samsun, Türkiye

Dr.Coşkun Gülser

Ondokuz Mayıs University, Faculty of Agriculture
Department of Soil Science and Plant Nutrition
55139 Samsun, Türkiye

Dr.Orhan Dengiz

Ondokuz Mayıs University, Faculty of Agriculture
Department of Soil Science and Plant Nutrition
55139 Samsun, Türkiye

Copyright © 2022 by Federation of Eurasian Soil Science Societies.

All rights reserved

ISBN

This Abstract book has been prepared from different abstracts sent to the symposium secretary only by making some changes in the format. Scientific committee regret for any language and/or aim-scope.

All rights reserved. No parts of this publication may be reproduced, copied, transmitted, transcribed or stored in any form or by any means such as mechanical, electronic, magnetic, optical, chemical, manual or otherwise, without prior written permission from copyright owner.

Publication date : 2 December 2022



Visit the Symposium web site at
<http://www.fesss.org/>

E-mail: symposium@fesss.org



With the support of the
Erasmus + Programme
of the European Union

Feasibility of re-involvement of the fallow soddy-podzolic soil in cultivation

Andrey LITVINOVICH ^{1,2}, Anton LAVRISHCHEV ², Vladimir BURE ^{1,3}, Tara GRUJIĆ ⁴,
Elmira SALJNIKOV ^{4,5,*}

¹ Agrophysical Research Institute, Grazhdanskaya 14, 195220 St. Petersburg, Russia

² St. Petersburg State Agrarian University, Peterburgskoye 2, 196601 St. Petersburg, Russia

³ St. Petersburg State University, Universitetskaya 7/9, 199034 St. Petersburg, Russia

⁴ Institute of Soil Science, Department of Pedology, Teodora Drajzera 7, 11000, Belgrade, Serbia

⁵ Mitscherlich Academy for Soil Fertility (MITAK) GmbH, Prof.-Mitscherlich-Allee 1, 14641 Paulinenaue, Germany

ABSTRACT

Transformations of the soddy-podzolic gleyic clay soil in a long-term agricultural use (> 200 years) was studied in Leningrad region, northwest Russia. This feasibility study investigated the possibility of re-cultivation of these soils after long term fallowing. Morphological structure, particle size distribution, content and ratio of ferrous and oxide forms of iron in the profile of virgin (indigenous forest) and arable drained soil were analysed. In addition, changes in the organogenic-profile were traced in the course of long-term agrogenesis (>200 years). In virgin forest soil, during its pedogenesis the loss of fractions <0.01 mm from the eluvial layer was 877.4 kg m⁻², and the loss of <0.0001mm was 287.5 kg m⁻², as compared with parent material not affected by the processes of pedogenesis. However, long-term agrogenesis (>200 years) led to increased eluvial losses of fine earth particles. The loss of fraction <0.01 mm from the arable horizons was 1244.8, and < 0.0001 mm was 570 kg m⁻², respectively. This was due to multiple yearly tillage that increased the porosity of the soil and thus intensified lessivage, which led to increased leaching and eluvial losses. The total loss of colloids from the entire profile of virgin soil was 262.1 kg m⁻², and from the arable layer of drained soil - 290.1 kg m⁻². The humus enrichment of the colloids of the plough (P) horizon of the arable soil was two times lower than that of the surface (AY) horizon of the virgin soil. The relative share of the participation of colloids in the fixation of humus by the soil was the same (11.6 and 10.9%, respectively). In the subsurface horizons, the absolute content of humus in the colloids decreased, and the share of participation in the fixation of humus increased. When soddy-podzolic gleyic clay soil is brought to cultivation then the water-air regime is improved, content and composition of humus, depth of arable horizon is increased and the soil acidity decreased. At the same time the leaching of fine earth materials is accelerated. When this soil was withdrawn from crop production, the positive changes achieved as a result of cultivation were gradually lost. For the first time we could qualitatively calculate the losses of the fine earth fractions for the given soil from top soil. Taking into account the high costs of re-cultivation of the former land and a high cost of re-installation and maintenance of an optimal hydrological regime (drainage network) we concluded that repeated ploughing and involvement of arable soddy-podzolic gleyic clay soil into cultivation is economically unreasonable

Key words:

Corresponding author : Elmira SALJNIKOV

E-mail: soils.saljnikov@gmail.com