



# **PROCEEDINGS OF THE**

# 30<sup>th</sup> International Symposium on Analytical and Environmental Problems

Szeged, Hungary October 7-8, 2024



# **University of Szeged**

# **Edited by:**

Tünde Alapi Róbert Berkecz István Ilisz

**Publisher:** University of Szeged, H-6720 Szeged, Dugonics tér 13, Hungary

ISBN 978-963-688-009-5

# 2024. Szeged, Hungary

## The 30<sup>th</sup> International Symposium on Analytical and Environmental Problems

### **Organized by:**

SZAB Kémiai Szakbizottság Analitikai és Környezetvédelmi Munkabizottsága

### **Supporting Organizations**

Institute of Pharmaceutical Analysis, University of Szeged Department of Inorganic and Analytical Chemistry, University of Szeged

## Symposium Chairman:

István Ilisz, DSc

### **Honorary Chairman:**

Zoltán Galbács, PhD

### **Organizing Committee:**

István Ilisz, DSc professor of chemistry University of Szeged, Institute of Pharmaceutical Analysis Tünde Alapi, PhD assistant professor University of Szeged, Department of Molecular and Analytical Chemistry Róbert Berkecz, PhD associate professor University of Szeged, Institute of Pharmaceutical Analysis

### **Scientific Committee:**

István Ilisz, DSc Tünde Alapi, PhD Róbert Berkecz, PhD Daniela Sojic Merkulov, PhD full professor University of Novi Sad, Faculty of Sciences, Department of Chemistry, Biochemistry and Environmental Protection

#### YIELD, MORPHOLOGICAL AND PHYSIOLOGICAL PARAMETERS OF ORGANIC AVENA SATIVA L. PLANT AS AFFECTED BY MINERAL, ORGANO-MINERAL AND STEEL SLAG AMENDMENTS TO ACID SOIL

#### <u>Aleksandra Stanojković-Sebić<sup>1</sup></u>, Dobrivoj Poštić<sup>2</sup>, Marina Jovković<sup>1</sup>, Radmila Pivić<sup>1</sup>

<sup>1</sup>Institute of Soil Science, T. Drajzera 7, 11000 Belgrade, Serbia <sup>2</sup>Institute for Plant Protection and Environment, T. Drajzera 9, 11000 Belgrade, Serbia e-mail: astanojkovic@yahoo.com

#### Abstract

Oat (Avena sativa L.) is one of the most important self-fertilizing field plant belonging to the Poaceae family, with multiple purposes as a fodder plant and in human nutrition. Of all the cereals, it tolerates heavy and wet soils the best, but with a very good reaction to fertilization [1]. The current study evaluated the effects of solely application of mineral (NPK) and organo-mineral (OM) fertilizers, and their individual combination with steel slag (SS), on the certain measured morphological parameters (mean number of leaves per plant - MNLP; minimum leaf length per plant - MinLL, in cm; maximum leaf length per plant - MaxLL, in cm; number of ears per plant - NEP), physiological parameters [content of: chlorophyll - Chl, flavonoids - Flv, anthocyanins -Ant, nitrogen balance index (Chl/Flav Ratio) - NBI], and the yield, in oat crops, between the treatments and in relation to the control. The experiment was performed in semi-controlled greenhouse conditions, in pots, from the 4th decade of March to the 4th decade of June, in 2024, using organic oat seeds, with Eutric Cambisol [2]. The soil is characterized as a light clay with an acid reaction. Steel slag was taken from the steel factory different deposition sites and characterized to have very alkaline reaction, Ca and Mg contents mainly in their oxide forms, and high total Fe and Mn content (but with lower amounts in their soluble forms) [3]. NBI and the contents of Chl, Flv, Ant, were measured using portable Dualex optical leafclip sensor (FORCE-A, Orsay, France). The obtained results showed that the MNLP, MinLL, MaxLL, NEP, the contents of Chl, Flv, Ant and NB index, in organic oats, significantly differed between the treatments applied and in relation to control. The effect of treatments on obtained oat yield insignificantly differed between the treatments, but significantly in relation to the control. The best results were obtained with the use of OM+SS, whereby good results were obtained with the NPK+SS use. Concluding, applied steel slag, in combination with mineral and organo-mineral fertilizer, respectively, showed the positive effects on the morphological and physiological parameters in organic oat and its yield, grown on acid type of soil. Regarding this research, in such combinations steel slag showed high potential in usage toward improvement of tested soil fertility without adverse effects.

#### Acknowledgements

Ministry of Science, Technological Development and Innovation of the Republic of Serbia, Contracts No. 451-03-66/2024-03/200011 and 451-03-66/2024-03/ 200010.

#### References

[1] Dj.Glamočlija, Field Production - Grains and Grain Legumes, Faculty of Agriculture, Zemun - Belgrade, Serbia, 2012, pp. 87. (in Serbian)

[2] IUSS Working Group WRB, World Reference Base for Soil Resources, 4th ed., Vienna, Austria, 2022.

[3] R.Pivić, A.Stanojković, S.Maksimović, D.Stevanović, Fresenius Env. Bull. 20 (2011) 875.