

ISSN 1821-1046

UDK 630

**INSTITUTE OF FORESTRY
BELGRADE**




**INSTITUT ZA ŠUMARSTVO
BEOGRAD**

SUSTAINABLE FORESTRY

ODRŽIVO ŠUMARSTVO

**COLLECTION
Vol. 89-90**

**ZBORNİK RADOVA
Vol. 89-90**



**BELGRADE BEOGRAD
2024.**

ISSN 1821-1046



9 771821 104000

ISSN 1821-1046
UDK 630

INSTITUTE OF FORESTRY
BELGRADE



INSTITUT ZA ŠUMARSTVO
BEOGRAD

SUSTAINABLE FORESTRY

COLLECTION
Vol. 89-90

ODRŽIVO ŠUMARSTVO

ZBORNIK RADOVA
Vol. 89-90

BELGRADE BEOGRAD
2024.

**INSTITUTE OF FORESTRY
BELGRADE
COLLECTION OF PAPERS**

**INSTITUT ZA ŠUMARSTVO
BEOGRAD
ZBORNİK RADOVA**

Publisher	Izdavač
Institute of Forestry Belgrade, Serbia	Institut za šumarstvo Beograd, Srbija
For Publisher	Za izdavača
Ljubinko Rakonjac, Ph.D.	Dr Ljubinko Rakonjac
Editor-in-Chief	Glavni i odgovorni urednik
Tatjana Ćirković-Mitrović, Ph.D.	Dr Tatjana Ćirković-Mitrović

Editorial Board

Redakcioni odbor

Ljubinko Rakonjac, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Ljubinko Rakonjac Institut za šumarstvo, Beograd, Srbija
Mara Tabaković-Tošić, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Mara Tabaković-Tošić Institut za šumarstvo, Beograd, Srbija
Biljana Nikolić, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Biljana Nikolić Institut za šumarstvo, Beograd, Srbija
Zoran Miletić, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Zoran Miletić Institut za šumarstvo, Beograd, Srbija
Milorad Veselinović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Milorad Veselinović Institut za šumarstvo, Beograd, Srbija
Aleksandar Lučić, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Aleksandar Lučić Institut za šumarstvo, Beograd, Srbija
Vladan Popović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Vladan Popović Institut za šumarstvo, Beograd, Srbija
Zlatan Radulović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Zlatan Radulović Institut za šumarstvo, Beograd, Srbija
Ljiljana Brašanac-Bosanac, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Ljiljana Brašanac-Bosanac Institut za šumarstvo, Beograd, Srbija
Saša Eremija, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Saša Eremija Institut za šumarstvo, Beograd, Srbija
Miroslava Marković, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Miroslava Marković Institut za šumarstvo, Beograd, Srbija
Sonja Braunović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Sonja Braunović Institut za šumarstvo, Beograd, Srbija
Đorđe Jović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Đorđe Jović Institut za šumarstvo, Beograd, Srbija
Katarina Mladenović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Katarina Mladenović Institut za šumarstvo, Beograd, Srbija
Suzana Mitrović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Suzana Mitrović Institut za šumarstvo, Beograd, Srbija
Snežana Stajić, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Snežana Stajić Institut za šumarstvo, Beograd, Srbija
Nevena Čule, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Nevena Čule Institut za šumarstvo, Beograd, Srbija
Ilija Đorđević, Ph.D. Institute of Forestry, Belgrade, Serbia	Ilija Đorđević, Ph.D. Institute of Forestry, Belgrade, Serbia
Goran Češljar, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Goran Češljar Institut za šumarstvo, Beograd, Srbija
Tomislav Stefanović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Tomislav Stefanović Institut za šumarstvo, Beograd, Srbija

Zoran Poduška, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Zoran Poduška Institut za šumarstvo, Beograd, Srbija
Tatjana Dimitrijević, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Tatjana Dimitrijević Institut za šumarstvo, Beograd, Srbija
Filip Jovanović, Ph.D. Institute of Forestry, Belgrade, Serbia	Dr Filip Jovanović Institut za šumarstvo, Beograd, Srbija
Assoc. Prof. Iantcho Naidenov, Ph.D. Forest Protection Station, Sofia, Bulgaria	Assoc. Prof. Dr Iantcho Naidenov Forest Protection Station, Sofia, Bulgaria
Prof. dr Makedonka Stojanovska, Faculty of Forestry, Ss. Cyril and Methodius University in Skopje, N. Macedonia	Prof. dr Makedonka Stojanovska, Šumarski fakultet Univerzitet Sv. Ćirilija i Metodija u Skoplju, S. Makedonija
Dr Zuzana Sarvašová National Forest Centre – Forest Research Institute, Slovakia	Dr Zuzana Sarvašová National Forest Centre – Forest Research Institute, Slovakia
Dr Alessandro Paletto Council for Agricultural Research and Economics, Italy	Dr Alessandro Paletto Savet za poljoprivredna istraživanja i ekonomiju, Italija
Associate Professor dr Sonia Quiroga Department of Economics, University of Alcalá, Spain	Associate Professor dr Sonia Quiroga Katedra za ekonomiju, Univerzitet u Alkali, Španija
Prof. dr Marijana Kapović Solomun Faculty of Forestry, Banja Luka, Republic of Srpska, Bosnia and Herzegovina	Prof. dr Marijana Kapović Solomun Šumarski fakultet, Banja Luka, Republika Srpska, Bosna i Hercegovina
Ph.D. Vanja Daničić Faculty of Forestry, Banja Luka, Republic of Srpska, Bosnia and Herzegovina	Dr Vanja Daničić Šumarski fakultet, Banja Luka, Republika Srpska, Bosna i Hercegovina
Dr.Sc. Mirza Dautbašić Faculty of Forestry, Sarajevo, Bosnia and Herzegovina	Dr Mirza Dautbašić Šumarski fakultet, Sarajevo, Bosna i Hercegovina
Dr. Sc. Muhamed Bajrić Faculty of Forestry, Sarajevo, Bosnia and Herzegovina	Dr Muhamed Bajrić Šumarski fakultet, Sarajevo, Bosna i Hercegovina
Dr. Sc. Alma Bogunić Hajrudinović Faculty of Forestry, Sarajevo, Bosnia and Herzegovina	Dr Alma Bogunić Hajrudinović Šumarski fakultet, Sarajevo, Bosna i Hercegovina
Doc.dr Milić Čurović Biotechnical Faculty, University of Montenegro, Montenegro	Doc. dr Milić Čurović Biotehnički fakultet, Univerzitet Crne Gore, Crna Gora
Assistant Professor dr Špela Pezdevšek Malovrh Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia	Assistant Professor dr Špela Pezdevšek Malovrh Biotehnički fakultet, Univerzitet Ljubljana, Ljubljana, Slovenija
Dr Dijana Vuletić Croatian Forest Research Institute, Jastrebarsko, Croatia	Dr Dijana Vuletić Hrvatski šumarski institut, Jastrebarsko, Hrvatska

Technical Editor and Layout

Ljiljana Brašanac-Bosanac, Ph.D.

Tehnički urednik i prelom teksta

Dr Ljiljana Brašanac-Bosanac

Secretary

M.Sc. Jelena Božović

Sekretar Zbornika

Mst. Jelena Božović

Printed in

100 copies

Tiraž

100 primeraka

Printed by

Black and White
Belgrade

Štampa

Black and White
Beograd

All rights reserved. No part of this publication might be reproduced by any means: electronic, mechanical, copying or otherwise, without prior written permission of the publisher.

Belgrade, 2024

Preuzimanje članaka ili pojedinih delova ove publikacije u bilo kom obliku nije dozvoljeno bez odobrenja izdavača.

Beograd, 2024

Cover Page: Author of the Photos, B.Sc. Nenad Šurjanac

Naslovna strana: Autor fotografije Nenad Šurjanac, dipl. inž.

CONTENT SADRŽAJ

Vol. 89-90

*Ivona KERKEZ JANKOVIĆ, Dragica VILOTIĆ, Marina NONIĆ,
Filip MAKSIMOVIĆ, Mirjana ŠJAJČIĆ-NIKOLIĆ*

**GENEPOOL OF WOODY SPECIES IN THE STRICT NATURE RESERVE
"FELJEŠANA"**

1

*Vladan POPOVIĆ, Aleksandar LUČIĆ, Aleksandar VEMIĆ, Sanja JOVANOVIĆ,
Ivona KERKEZ-JANKOVIĆ, Mirjana ŠJAJČIĆ-NIKOLIĆ*

**WHITE WILLOW (SALIX ALBA L.) VARIABILITY IN THE
LANDSCAPES OF OUTSTANDING FEATURES "GREAT WAR ISLAND"
BASED ON MORPHOLOGICAL TRAITS OF THE LEAVES: A BASIS
FOR ASSESSMENT OF GENE POOL**

17

Alen GAČIĆ, Marijana KAPOVIĆ SOLOMUN, Ilija ČIGOJA, Saša EREMIJA
**CHARACTERISTICS OF SOILS IN FOREST MANAGEMENT UNIT
„MALA UKRINA“**

31

*Snežana STAJIĆ, Vlado ČOKEŠA, Ljubinko RAKONJAC, Saša EREMIJA,
Suzana MITROVIĆ, Zoran PODUŠKA, Branka PAVLOVIĆ*

**PHYTOCOENOLOGICAL ANALYSIS OF SESSILE OAK AND TURKEY
OAK FORESTS (*QUERCETUM PETRAEAE-CERRIDIS* B. JOVANOVIĆ
1979, S.L.) IN THE TERRITORY OF KOSMAJ**

47

*Branka PAVLOVIĆ, Vlado ČOKEŠA, Snežana STAJIĆ, Violeta BABIĆ,
Zoran PODUŠKA, Nikola MARTAĆ, Branko KANJEVAC*

**PLANT SPECIES AS HABITAT INDICATORS IN BEECH FORESTS
FOLLOWING CLEARCUTTING**

63

*Snežana OBRADOVIĆ, Milan MEDAREVIĆ, Damjan PANTIĆ, Biljana ŠLJUKIĆ,
Nenad PETROVIĆ, Dragan BOROTA, Aleksandar POPOVIĆ*

**SPONTANEOUS DEVELOPMENT OF MIXED STANDS OF FIR, SPRUCE
AND BEECH ON MT. TARA**

77

*Suzana MITROVIĆ, Milorad VESELINOVIC, Snežana STAJIĆ,
Renata GAGIĆ-SERDAR, Miroslava MARKOVIĆ, Ivana BJEDOV,
Marija MILOSAVLJEVIĆ*

**EFFECTS OF FERTILISATION ON SURVIVAL AND
MORPHOLOGICAL GROWTH CHARACTERISTICS OF ONE-YEAR-
OLD SEEDLINGS OF PAULOWNIA ELONGATA S.Y. HU. AND
PAULOWNIA FORTUNEI SEEM. HEMS. IN TWO DIFFERENT SITES
IN SERBIA**

87

<i>Filip JOVANOVIĆ, Ivana ŽIVANOVIĆ, Nenad ŠURJANAC, Đorđe FILIPOVIĆ, Đorđe JOVIĆ, Aleksandar LUČIĆ</i> CONDITION OF DOUGLAS FIR TREES IN THE URBAN AREA OF BELGRADE (SERBIA)	109
<i>Aleksandar VEMIĆ, Zlatan RADULOVIĆ, Katarina MLADENović, Ljubinko RAKONJAC</i> THE MOST COMMON FUNGI ASSOCIATED WITH A DECLINE OF TURKEY OAK (<i>QUERCUS CERRIS</i> L.) IN URBAN CONDITIONS IN SERBIA	119
<i>Katarina MLADENović, Aleksandar VEMIĆ, Sabahudin HADROVIĆ, Milan KABILJO, Đorđe JOVIĆ</i> A CONTRIBUTION TO THE KNOWLEDGE OF THE MITES (ACARI) FAUNA OF THE HORNBEAM IN SERBIA	131
<i>Miroslava MARKOVIĆ, Renata GAGIĆ-SERDAR, Goran ČEŠLJAR, Suzana MITROVIĆ, Đorđe JOVIĆ, Mihajlo MARKOVIĆ</i> USE OF A DATABASE FOR DETERMINING THE SPATIAL DISTRIBUTION OF PESTS AND DISEASES IN THE FORESTS OF SERBIA	141
<i>Jelena BOŽOVIĆ, Zlatan RADULOVIĆ, Bojan KONATAR, Snežana STAJIĆ, Nevena ČULE, Radojica PIŽURICA, Dragana ŽIVOJINOVIĆ</i> ANALYSIS OF THE CHEMICAL COMPOSITION OF THREE FUNGAL SPECIES WITH MEDICINAL PROPERTIES TO INVESTIGATE THEIR MEDICAL AND ECOLOGICAL POTENTIAL	149
<i>Marija S. MARKOVIĆ, Biljana M. NIKOLIĆ, Dejan S. PLJEVLJAKUŠIĆ, Ljubinko B. RAKONJAC, Sonja Z. BRAUNOVIĆ, Filip A. JOVANOVIĆ, Vesna P. STANKOV JOVANOVIĆ</i> TRADITIONAL MEDICINAL USE OF PLANTS FROM THE GENUS CRATAEGUS IN THE PIROT DISTRICT (SERBIA)	161
<i>Olga GAJANIĆ, Biljana JOVIĆ, Ivana BJEDOV, Marija NEŠIĆ</i> THE POSSIBILITY OF CREATING AN EDUCATIONAL TRAIL INSPIRED BY THE MEDICINAL AND USEFUL PROPERTIES OF THE SHRUB SPECIES PRESENT IN THE ARBORETUM OF THE FACULTY OF FORESTRY	177
<i>Jelena UROŠEVIĆ, Dragica STANKOVIĆ, Goran TRIVAN, Đorđe JOVIĆ, Saša ORLOVIĆ, Sonja BRAUNOVIĆ, Filip JOVANOVIĆ</i> CO-FIRING OF CONTAMINATED WILLOW BIOMASS (<i>SALIX</i> L.) WITH LIGNITE IN THE ENERGY PRODUCTION PROCESS	199
<i>Ljiljana BRAŠANAC-BOSANAC, Nevena ČULE, Ilija ĐORĐEVIĆ, Goran ČEŠLJAR, Aleksandar LUČIĆ, Predrag ŠUMARAC, Tatjana ĆIRKOVIĆ-MITROVIĆ</i> THE IMPORTANCE OF APPLYING THE CIRCULAR BIOECONOMY CONCEPT IN FORESTRY	211

Goran ĐORĐEVIĆ, Martina PETKOVIĆ, Marko TOMIĆ, Andreja MIJATOVIĆ

**FOREST FIRES AS AN ECOLOGICAL SAFETY FACTOR AND ITS
IMPACT ON SUSTAINABLE DEVELOPMENT**

223

A GUIDE FOR WRITING RESEARCH PAPER

233

DOI: 10.5937/SustFor2490077O

UDK: 630*188(497.11Tara)=111

Original scientific paper

SPONTANEOUS DEVELOPMENT OF MIXED STANDS OF FIR, SPRUCE AND BEECH ON MT. TARA

Snežana OBRADOVIĆ¹, Milan MEDAREVIĆ¹, Damjan PANTIĆ¹,
Biljana ŠLJUKIĆ¹, Nenad PETROVIĆ¹, Dragan BOROTA¹,
Aleksandar POPOVIĆ¹*

Abstract: *The main objective of this study is to analyze the structural production and dynamic changes of mixed forests of fir, spruce and beech on Mt. Tara, in permanent sample plots which were exempt from the regular management over a period of more than 60 years (1955-2017). The results of the research indicate that during the analyzed period mixed forests of fir, spruce and beech have undergone a number of changes in structural and production terms. The significant presence of trees of large dimensions (volume accumulation) slowed down the developmental dynamics of these stands and made regeneration and recruitment difficult, which had a negative impact on their structural development and, in general, on the stability of these stands. The results show that selection forests are possible to achieve only with a permanent management approach and adherence to the modern principles of close-to-nature forest management. In this sense, selection cutting in the context of close-to-nature forest management is imposed as a means of accelerating the dynamics of development of these forests, repairing their structural development (by achieving and preserving their selection structure), ecological stability and functional values. On the contrary, the absence of activity and spontaneous development of these stands leads to a general devitalization and a deviation from the desired goals.*

Keywords: Mt. Tara, forests of fir, spruce and beech, spontaneous stand development, stand structure, stand dynamics

SPONTANI RAZVOJ MEŠOVITIH SASTOJINA JELE, SMRČE I BUKVE NA TARI

Sažetak: *Osnovni cilj ovih istraživanja jeste da se analiziraju strukturne, proizvodne i dinamičke promene mešovitih šuma jele, smrče i bukve na Tari, na stalnim oglednim površinama, koje su izuzete iz redovnog gazdovanja u periodu dužem od 60 godina (1955-2017). Rezultati istraživanja ukazuju na to da su u analiziranom periodu mešovite šume jele, smrče i bukve pretrpele niz promena u strukturnom i proizvodnom smislu. Značajno prisustvo stabala jakih dimenzija (nagomilavanje zapremine) usporilo je dinamiku razvoja ovih sastojina i otežalo podmlađivanje i urastanje, što se negativno*

¹Faculty of forestry, Kneza Visaslava 1, 11.000 Belgrade, Serbia

*Corresponding author. E-mail: snezana.obradovic@sfb.bg.ac.rs

© 2024 The Authors. Published by the Institute of Forestry, Belgrade.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

odrazilo na njihovu strukturnu izgrađenost i, generalno, na stabilnost ovih sastojina. Rezultati pokazuju da je prebirna šuma moguća jedino uz permanentan gazdinski pristup i poštovanje savremenih principa prirodi bliskog gazdovanja. U tom smisli, prebirna seča, u kontekstu prirodi bliskog gazdovanja, nameće se kao sredstvo ubrzanja dinamike razvoja ovih šuma, popravke njihove strukturne izgrađenosti (postizanja i očuvanja prebirne strukture), ekološke stabilnosti i funkcionalne vrednosti. Suprotno, odsustvo aktivnog delovanja i spontani razvoj sastojina vode ka njihovoj devitalizaciji i udaljavanju od željenih ciljeva.

Ključne reči: Tara, šume jele, smrče i bukve, spontani razvoj, struktura sastojine, dinamika sastojine

1. INTRODUCTION

The forest of fir, spruce and beech *Piceo-Abietetum* Col. 1965 (syn. *Piceo-Fago Abietetum* Chol. 1965; *Piceo-Abieti-Fagetum moesiicum* Mišić et al., 1978) is a specific three-dominant community, which can be found in Serbia on Mt. Tara, Mt. Zlatar, Mt. Golija, the Mojstirsko-Draške mountains, Mt. Kopaonik, Mt. Stara planina, and Pešter Plateau, etc. In Mt. Tara and the Pešter Plateau, this community occurs as a climate regional belt (Gajić, 1992; Tomić, Rakonjac, 2013). The ratio of spruce, fir, and beech in the association changed during the historical development of the vegetation, depending on the climatic conditions and the way of management (Tomić, Rakonjac, 2013).

These forests are characterized by structural diversity, i.e. a series of transitions from diverse to complex selection structural forms. In recent decades, there has been a growing interest of the international forestry circles in selection (uneven-aged) forest management systems, as these systems have been recognized as more natural, associated with more pronounced diversity and a greater social value of forests (Schütz, 2002). Increasingly, the focus is placed on the management methods and systems that are in harmony with nature and at the same time oriented toward multifunctional use. It is precisely the selection forest that is a silvicultural form suitable for producing multiple effects. It can therefore be referred to as a system in which the interests are balanced and the use is regulated as a compromise.

The selection forest is not created naturally, but solely in a way that continuously supports this concept by silvicultural measures. The structural changes, productivity and dynamics of development of the selection forests of fir, spruce and beech on Mt. Tara have been monitored since 1955 in a series of permanent sample plots (SPs). These SPs were exempt from regular management and left to spontaneous development and self-regulatory processes. The aim was to identify and analyze certain laws and trends in the development of these forests, as well as to implement new knowledge into a management process that aims at structurally and ecologically stable and highly functional selection forests. In those terms, from a time distance of more than 60 years, the goal of these studies was defined, and the following hypotheses were formulated:

- in the analyzed period (1955-2017) there were structural disturbances of the stands,
- the developmental dynamics of the stands was slowed down and regeneration and recruitment became difficult,
- the stability and functional value of these forests were reduced.

2. MATERIAL AND METHODS

2.1 Research object

The research was carried out in the "Tara" MU, which is an integral part of the national park with the same name. From the series of permanent SPs in this locality, sample plots SP-1, SP-2 and SP-4 were selected, which are located in the type of spruce, fir and beech forest (*Piceo - Abieti - Fagetum typicum*), on deep to medium deep brown soils on limestone. They were established in 1955 and subsequently largely left to spontaneous development. The basic spatial features of the SPs are shown in Table 1.

Table 1. *The basic spatial features of the SPs*

SP	Area (ha)	Altitude (m)	Exposure	Terrain slope (°)	Coordinates WGS N	Coordinates WGS E
SP-1	1.44	1099-1116	East	10-15 °	43 ° 55'10 " N 43 ° 55'16 " N	19 ° 25 '06"E 19 ° 25'11"E
SP-2	1.04	1157-1174	Southeast	5- 10 °	43 ° 54'46"N 43 ° 54'51"N	19 ° 24'33"E 19 ° 24'40"E
SP-4	1.00	1152-1174	West	10-15 °	43 ° 54'43"N 43 ° 54'48"N	19 ° 25'42"E 19 ° 25'45"E

The average annual air temperature on Mt. Tara is 5.2°C, the coldest month being January, while the warmest is July, with an average temperature of 13.8°C. The average air temperature in the vegetation period is 10.8°C, while in autumn the mean temperature is 6.1°C, which makes it warmer than spring (4.6 ° C). Annual precipitation is 1,005 mm of water sediment. The rainiest month is May and January and March are the driest months. Rainfall is more abundant during the vegetation period (584 mm of water sediment), which favors the development of forest vegetation.

2.2 Data collection and processing

The surveys used data from periodic measurements (1955, 1975, 2005 and 2017) of permanent SPs in the Mt. Tara forests, which are in the database of the Chair of Forest Management Planning of the Faculty of Forestry in Belgrade. The SPs were spatially defined by marking their boundaries and determining their coordinates, altitude, slope and exposure, while each tree in them was numbered. During periodic measurements, the boundaries of the SPs and the tree numbers were renewed, and the numbering of newly-recruited trees was also performed.

The measurement in the SPs involved a cross measurement of the diameter at breast height and a measurement of the height of each tree above the forest estimation threshold of 10 cm. The diameter was measured with an accuracy of 1 mm and height with an accuracy of 0.1 m.

To enable a reliable comparative analysis, the data of periodic measurements were processed using the same methodology. The volume was calculated by the volume tables method, using the local tables for fir and spruce on Mt. Tara (Banković, 1991) and general tables for high beech forests of Serbia (Mirković, 1969). The current volume increment method was used to calculate the current volume increment, whereby the increment percentage was determined by regression models expressing its dependence on the number of trees per unit area, the share of a particular tree species in the mixture and the diameter and height of the mean stand tree (Banković et al. 2000, 2002).

The diversity of tree dimensions in the SPs was estimated on the basis of the Gini coefficient (GCy) (Lexerød & Eid, 2006; Klopčič & Bončina, 2011), which was obtained using the formula:

$$GCy = \frac{\sum_{j=1}^n (2j - N - 1) \times g_j}{\sum_{j=1}^n g_j \times (N - 1)}$$

in which:

y- is the year in which the measurement was performed,

j- is the rank of trees ordered from 1 to n (from the thinnest to the thickest),

N- is the total number of trees,

g- is the basal area of a tree.

Statistical data processing was performed by applying descriptive statistics. Using the data of periodic measurements, the arithmetic mean, minimum, maximum and variation coefficient were calculated in each SP as the basic measures of variation in the number of trees (N), basal area (G), volume (V) and current volume increment (Iv) per hectare.

Within this chapter there can be subtitles of first and second line. In the description of material, it should be given enough information to allow other researchers to repeat the experiment at a different location. It is necessary to provide information on the material, subject of the study that precisely defines its origin, physical characteristics etc. If a device or instrument is used to obtain experimental results should be specified: name of the device or instrument, model, manufacturer's name and country of origin. If a scientifically recognized method is used it has to be cited in the References, without the explanation of the steps of the used method. If changes were made in a scientifically recognized method should be provided the original literature references that will support – justify those changes.

3. RESULTS AND DISCUSSION

The variation coefficient of the number of trees is 26.6% in SP-1, 19.3% in SP-2 and 33.6% in SP-4 (Table 2), indicating a more pronounced variation of this element over time. There is an evident decrease in the total number of trees compared to the beginning of the analyzed period and it amounts to 30.9% in SP-1, 32.1% in SP-2 and 50.5% in SP-4. The accumulation of trees of large dimensions ($d \geq 50$ cm) (Figure 1) resulted in the reduction of light in the stand, the reduction of growth space, as well as the absence of recruitment, which led to the extinction of thin trees (trees of lower storeys), i.e. to a decrease in the total number of trees in the observed period. Such a trend led to structural disturbances and tree distribution curves developed a flatter form, thus moving away from the shape that is typical of selection forests (Figure 1). The consequence of these changes is also a decrease in dimensional diversity, as indicated by the Gini coefficient, which declines over time (Figure 1). The trend of this coefficient in the observed period ranged from 0.556 to 0.547 in SP-1, i.e. from 0.557 to 0.536 in SP-2 and from 0.493 to 0.451 in SP-4.

Table 2. *Measures of variation in the basic numerical elements in the SPs*

SP	\bar{N} (trees · ha ⁻¹)	min.	max.	C _N (%)	\bar{G} (m ² · ha ⁻¹)	min.	max.	C _G (%)
SP-1	436	327	564	26.6	47.1	36.2	52.4	7.4
SP-2	550	422	650	19.3	47.5	35.2	55.7	9.9
SP-4	609	384	789	33.6	46.3	36.7	51.9	6.8
SP	\bar{V} (m ³ · ha ⁻¹)	min.	max.	C _V (%)	\bar{I}_V (m ³ · ha ⁻¹)	min.	max.	C _{I_V} (%)
SP-1	715	498	828	20.8	13.3	10.6	15.1	14.9
SP-2	701	450	898	31.6	14.1	10.2	17.3	22.4
SP-4	659	457	796	21.8	13.8	13.1	15.1	5.7

The basal area increased significantly in all SPs, i.e. by 44.3% in SP-1, 57.4% in SP-2 and by 26.4 % in SP-4 compared to the 1955 diameter. At the end of the period in 2017, the value of the basal area was high and it amounted to 52.4 m²·ha⁻¹ in SP-1, 55.4 m²·ha⁻¹ in SP-2 and 46.4 m²·ha⁻¹ in SP-4. The significant increase in the basal area can only be explained by the accumulation of trees of large dimensions.




Figure 1. Changes in the distribution of the number of trees by diameter classes and the Gini coefficient




Figure 2. Changes in the volume distribution by Biolay diameter classes

In 62 years' time the volume per hectare increased by 66.1% in SP-1 and 94.9% in SP-2 and by 52.2% in SP-4. The current volume values are high and range from $659.1 \text{ m}^3 \cdot \text{ha}^{-1}$ in SP-4 to $866.1 \text{ m}^3 \cdot \text{ha}^{-1}$ in SP-2. The volume structure by Biolay diameter classes also varied in favor of the category of trees of large dimensions. In all SPs, the share of thin and medium-sized trees in the total volume declined from one period to another, and the share of trees with diameters of over 50 cm increased (Figure 2). It is worrying that the share of thin trees in all SPs was lower than 10%, and according to Biolay, it should not be less than 20%, i.e. according to more recent research by Schütz (2001), lower than 15%.

The current volume increment increases over time. At the end of the analyzed period, the increase was high and it amounted to $14.4 \text{ m}^3 \cdot \text{ha}^{-1}$ in SP-1,

15.9 $\text{m}^3 \cdot \text{ha}^{-1}$ in SP-2 and 12.9 $\text{m}^3 \cdot \text{ha}^{-1}$ in SP-4. The aforementioned amount of increase indicates a still high production of the stands, but the pronounced disturbances regarding the reduction of the share of thinner, and as a rule, more vital and trees with a more intensive increment, indicate that this trend of current volume increment will not last long (Obradović, 2015).

4. CONCLUSION

The starting hypotheses of this research have been confirmed. Structural disturbances of fir, spruce and beech forests in the analyzed SPs in Mt. Tara were found. They arose as a result of the decrease in the total number of trees, which also resulted in a decrease in dimensional diversity. The decrease of the share of thin trees is particularly pronounced, as well as the increase in the share of trees of large dimensions ($d \geq 50$ cm). The distribution curves have developed a flatter form and are moving further away from the distribution that is typical of the selection forest.

Despite the reduction in the total number of trees, the accumulation of trees of large dimensions resulted in high values of basal area and volume per hectare. High absolute values of the current volume increment were also noted, which all together indicate a good production potential of a particular site. However, high volume does not always mean the desired state, especially if it is distributed among old trees (of large dimensions), of reduced technical value and if it slows down the development of the stand.

Insufficient regeneration and recruitment are the consequence of the stated distribution of trees by diameter classes, that is, reduced space for growth, with a negative impact on the preservation of these stands as stable and functional forms.

The trends observed in the fir, spruce and beech forests of Mt. Tara indicate that it is necessary to conduct permanent monitoring of all developmental processes, implement appropriate silvicultural and management measures that regulate them, and over time adjust and direct them towards the set goal, which is a structurally stable, socio-economically and ecologically valuable selection forest as a sustainable category. The leaving of stands to spontaneous development and self-regulation processes that happen over time, as is the case in the investigated SPs, moves us further away from this goal.

Acknowledgement: *We thank the Ministry of Science, Technological Development and Innovation of the Republic of Serbia, which financed this research based on the Agreement on the implementation and financing of scientific research work of the National Institute of Technology in 2024, registration number 451-03-65/2024-03/ 200169 dated 05.02.2024. year*

REFERENCES

Banković, S. (1991): Zapreminske tablice za jelu (*Abies alba* Mill.) i smrču (*Picea excelsa* Link.) u šumama nacionalnog parka Tara, Univerzitet u Beogradu-Šumarski fakultet, Nacionalni park Tara, Bajina Bašta (106).

Banković, S., Jović, D., Medarević, M., Pantić D. (2000): Regresioni modeli procenta zapreminskog prirasta u čistim i mešovitim sastojinama bukve i hrasta kitnjaka u Srbiji. Glasnik Šumarskog fakulteta 83, Univerzitet u Beogradu – Šumarski fakultet, Beograd (21-31).

Banković, S., Medarević, M., Pantić, D. (2002): Regresioni modeli procenta zapreminskog prirasta u najzastupljenijim sastojinama četinarskih vrsta drveća u Srbiji. Glasnik Šumarskog fakulteta 85, Univerzitet u Beogradu-Šumarski fakultet, Beograd (25-35).

Gajić, M., Kojić, M., Karadžić, D., Vasiljević, M., Stanić, M. (1992): Vegetacija Nacionalnog parka Tara, Šumarski fakultet, Beograd, Nacionalni park Tara, Bajina Bašta.

Klopčič, M., Bončina, A. (2011): Stand dynamics of silver fir (*Abies alba* Mill.) - European beech (*Fagus sylvatica* L.) forests during the past century: a decline of silver fir? Forestry 84 (3) (259–271).

Lexerød, N.L., Eid, T. (2006): An evaluation of different diameter diversity indices based on criteria related to forest management planning. Forest Ecology and Management 222 (17–28).

Mirković, D. (1969): Priručnik za određivanje zapremine i zapreminskog prirasta u bukovim sastojinama SR Srbije pri uređajnim radovima, Institut za šumarstvo i drvnu industriju, Beograd u Nikolić S., Banković S. (1992): Tablice i tehničke norme u šumartstvu, Zavod za udžbenike i nastavna sredstva, Beograd (1-257).

Obradović, S. (2015): Stanje i razvoj sastojina bukve, jele i smrče prašumskog porekla u Srbiji kao osnov za planiranje i obezbeđivanje prirodi bliskog gazdovanja. Ph.D. Thesis, University of Belgrade, Belgrade.

Schütz, J.-Ph. (2001): Der Plenterwald und weitere Formen strukturierter und gemischter Wälder. Parey, Berlin.

Schütz, J.-Ph. (2002): Uneven-aged silviculture: tradition and practices Forestry 75(4) (327-328).

Tomić, Z., Rakonjac, Lj. (2013): Šumske fitocenoze Srbije, priručnik za šumare, ekologe i biologe, Univerzitet Singidunum –Fakultet za primenjenu ekologiju, Institut za šumarstvo, Beograd (177).

SPONTANEOUS DEVELOPMENT OF MIXED STANDS OF FIR, SPRUCE AND BEECH FROM PERMANENT SAMPLE PLOTS ON MT. TARA

Snežana OBRADOVIĆ, Milan MEDAREVIĆ, Damjan PANTIĆ, Biljana ŠLJUKIĆ,
Nenad PETROVIĆ, Dragan BOROTA, Aleksandar POPOVIĆ

Summary

The forest of fir, spruce and beech *Piceo-Abietetum* Col. 1965 (syn. *Piceo-Fago Abietetum* Chol. 1965; *Piceo-Abieti-Fagetum moesiicum* Mišić et al., 1978) is a specific three-dominant community, which can be found in Serbia on Mt. Tara, Mt. Zlatar, Mt.

Golija, the Mojstirsko-Draške mountains, Mt. Kopaonik, Mt. Stara planina, and Pešter Plateau, etc. These forests are characterized by structural diversity, i.e. a series of transitions from diverse to complex selection structural forms. The structural changes, productivity and dynamics of development of the selection forests of fir, spruce and beech on Mt. Tara have been monitored since 1955 in a series of permanent sample plots (SPs). These SPs were exempt from regular management and left to spontaneous development and self-regulatory processes. The surveys used data from periodic measurements (1955, 1975, 2005 and 2017) of permanent SPs in the Mt. Tara forests, which are in the database of the Chair of Forest Management Planning of the Faculty of Forestry in Belgrade. The results of the research indicate that during the analyzed period mixed forests of fir, spruce and beech have undergone a number of changes in structural and production terms. The significant presence of trees of large dimensions (volume accumulation) slowed down the developmental dynamics of these stands and made regeneration and recruitment difficult, which had a negative impact on their structural development and, in general, on the stability of these stands. The results show that selection forests are possible to achieve only with a permanent management approach and adherence to the modern principles of close-to-nature forest management. In this sense, selection cutting in the context of close-to-nature forest management is imposed as a means of accelerating the dynamics of development of these forests, repairing their structural development (by achieving and preserving their selection structure), ecological stability and functional values. On the contrary, the absence of activity and spontaneous development of these stands leads to a general devitalization and a deviation from the desired goals.

SPONTANI RAZVOJ MEŠOVITIH SASTOJINA JELE, SMRČE I BUKVE SA STALNIH OGLEDNIH POVRŠINAMA NA TARI

Snežana OBRADOVIĆ, Milan MEDAREVIĆ, Damjan PANTIĆ, Biljana ŠLJUKIĆ,
Nenad PETROVIĆ, Dragan BOROTA, Aleksandar POPOVIĆ

Rezime

Šuma jele, smrče i bukve *Piceo-Abietetum* Čol. 1965. (syn. *Piceo-Fago Abietetum* Čol. 1965.; *Piceo-Abieti-Fagetum moesiaticum* Mišić *et al.*, 1978.) je specifična trodominantna zajednica, koja je u Srbiji zastupljena na Tari, Zlataru, Goliji, Mojstirsko-Draškim planinama, Kopaoniku, Staroj planini, Pešterskoj visoravni itd. Ove šume se karakterišu strukturnom raznolikosti, tj. nizom prelaza od raznodobnih ka prebirnim strukturnim oblicima. Strukturne promene, proizvodnost i dinamika razvoja prebirnih šuma jele, smrče i bukve na Tari prate se od 1955. godine, na seriji stalnih oglednih površina (OP). Ove OP su izuzete iz redovnog gazdovanja i prepuštene su spontanom razvoju i samoregulacijskim procesima. U istraživanjima su korišćeni podaci periodičnih premera (1955, 1975, 2005. i 2017. godine) stalnih OP u šumama Tare, koji se nalaze u bazi podataka Katedre Planiranja gazdovanja šumama, Šumarskog fakulteta u Beogradu. Rezultati istraživanja ukazuju na to da su u analiziranom periodu mešovite šume jele, smrče i bukve pretrpele niz promena u strukturnom i proizvodnom smislu. Značajno prisustvo stabala jakih dimenzija (nagomilavanje zapremine) usporilo je dinamiku razvoja ovih sastojina i otežalo podmlađivanje i urastanje, što se negativno odrazilo na njihovu strukturnu izgrađenost i, generalno, na stabilnost ovih sastojina. Rezultati pokazuju da je prebirna šuma moguća jedino uz permanentan gazdinski pristup i poštovanje savremenih principa prirodi bliskog gazdovanja. U tom smislu, prebirna seča, u kontekstu prirodi bliskog gazdovanja, nameće se kao sredstvo ubrzanja dinamike razvoja ovih šuma, popravke njihove strukturne izgrađenosti (postizanja i očuvanja prebirne strukture),

ekološke stabilnosti i funkcionalne vrednosti. Suprotno, odsustvo aktivnog delovanja i spontani razvoj sastojina vode ka sveopštoj devitalizaciji i udaljavanju od željenih ciljeva.

CIP - Каталогизација у публикацији
Народна библиотека Србије, Београд

630

SUSTAINABLE Forestry : collection =
Održivo šumarstvo = zbornik radova /
editor-in-chief Tatjana Ćirković-Mitrović. -
2008, t. 57/58- . - Belgrade: Institute of
forestry, 2008- (Beograd : Black and
White). - 24 cm

Godišnje. - Je nastavak: Zbornik radova -
Institut za šumarstvo = ISSN 0354-1894
ISSN 1821-1046 = Sustainable Forestry
COBISS.SR-ID 157148172